



Physiological, metabolic and biotechnological features of extremely thermophilic microorganisms

James A. Counts,¹ Benjamin M. Zeldes,¹ Laura L. Lee,¹ Christopher T. Straub,¹ Michael W.W. Adams² and Robert M. Kelly^{1*}

The current upper thermal limit for life as we know it is approximately 120°C. Microorganisms that grow optimally at temperatures of 75°C and above are usually referred to as 'extreme thermophiles' and include both bacteria and archaea. For over a century, there has been great scientific curiosity in the basic tenets that support life in thermal biotopes on earth and potentially on other solar bodies. Extreme thermophiles can be aerobes, anaerobes, autotrophs, heterotrophs, or chemolithotrophs, and are found in diverse environments including shallow marine fissures, deep sea hydrothermal vents, terrestrial hot springs—basically, anywhere there is hot water. Initial efforts to study extreme thermophiles faced challenges with their isolation from difficult to access locales, problems with their cultivation in laboratories, and lack of molecular tools. Fortunately, because of their relatively small genomes, many extreme thermophiles were among the first organisms to be sequenced, thereby opening up the application of systems biology-based methods to probe their unique physiological, metabolic and biotechnological features. The bacterial genera *Caldicellulosiruptor*, *Thermotoga* and *Thermus*, and the archaea belonging to the orders Thermococcales and Sulfolobales, are among the most studied extreme thermophiles to date. The recent emergence of genetic tools for many of these organisms provides the opportunity to move beyond basic discovery and manipulation to biotechnologically relevant applications of metabolic engineering. © 2017 Wiley Periodicals, Inc.

How to cite this article:

WIREs Syst Biol Med 2017, e1377. doi: 10.1002/wsbm.1377

INTRODUCTION

Extreme thermophiles are distinct from other organisms due to their ability to subsist optimally at temperatures in excess of 75°C. Their survival in these harsh environments piqued the interest of

curious microbiologists as far back as the turn of the 20th century. In fact, one of the earliest reports of thermophiles occurred in 1903 describing bacterial samples taken from pools in Yellowstone National Park.¹ Although this drew interest and debate about the limits of life and our evolutionary history, the study of thermophiles did not begin in earnest until the 1960s. Around this time, extensive sampling projects in Yellowstone lead to the isolation of *Thermus aquaticus*² (an aerobic bacterium with optimal growth between 70 and 75°C), known for its DNA polymerase that revolutionized the field of molecular biology through its use in the polymerase chain reaction (PCR). This enzyme in particular represented

*Correspondence to: rmkelly@ncsu.edu

¹Department of Chemical and Biomolecular Engineering, North Carolina State University, Raleigh, NC, USA

²Department of Biochemistry and Molecular Biology, University of Georgia, Athens, GA, USA

Conflict of interest: The authors have declared no conflicts of interest for this article.

one of the earliest uses of thermally stable enzymes for a biotechnological application. The next few decades yielded the discovery of thermophiles around the globe in extremely diverse environments, ranging from volcanoes and calderas to deep sea smoker vents to terrestrial mud pools.^{3–7}

The apparent diversity and novelty of these microbes likely drove early research in this field to uncover the molecular machinery central to their survival. In fact, some of the earliest sequenced genomes were extremophiles,^{8–10} furthering efforts to understand the molecular and genetic basis for thermophily and the evolution of life. However, a lack of genetics tools has impeded the extensive study of these organisms by traditional approaches (i.e., gene deletions to understand the consequences of loss of function). In lieu of more traditional methods, the availability of genomic data for many extreme thermophiles supported 'omics'-based approaches to ascertain the function of specific genes and their roles in the unique biochemistry of these organisms. As such, the merger of systems biology (e.g., transcriptomics and genomics), traditional microbiological studies, and newly emerging genetic systems¹¹ is opening the door for metabolic engineering opportunities to bring extreme thermophiles into the technological limelight. This will allow for these organisms to be utilized as sources of uniquely functioning enzymes, optimized niche industrial strains, and novel metabolic engineering platforms. Such opportunities for biotechnological application are already being pursued for members of the bacterial genera *Caldicellulosiruptor* and *Thermotoga* and for archaea belonging to the orders Thermococcales and Sulfolobales. Here we present a brief overview of these extremely thermophilic organisms, with the intention of highlighting potential biotechnological applications, which exploit their distinctive metabolisms.

SULFOLOBALES

Perhaps the most distinctive subject matter for this review focuses on the extreme thermoacidophiles from order Sulfolobales. The Sulfolobales comprise an order of archaea taxonomically defined within the class Thermoprotei, within the phylum Crenarchaeota. These organisms inhabit environments characterized by both extreme temperatures (65–90°C) and low pH (1.0–3.5), such as terrestrial solfatara or mud pools, often closely associated with volcanic activity and laden with inorganic materials.¹² In fact, the first species of the order to be isolated, *Sulfolobus acidocaldarius* (from Locomotive Spring in Yellowstone National Park), was reported to oxidize sulfur

to fuel autotrophic growth, leading to the name Sulfolobales.⁷ However, this phenotype has not been observed in the currently studied *S. acidocaldarius* type strains, although many isolates from the genera *Sulfolobus*, *Metallosphaera* and *Acidianus* utilize S⁰ as an electron donor.^{5,13–21} Thus, reports that *S. acidocaldarius* strains from culture collections cannot⁵ oxidize S⁰ suggests that repeated passages on rich media have led to the loss of this ability or that inherent difficulties exist in isolating pure cultures from environmental enrichments. Beyond sulfur oxidation, several species, especially those from the genus *Acidianus*, *Sulfurisphaera*, and *Stygiolobus*, are capable of sulfur reduction, and often utilize hydrogen to produce hydrogen sulfide as a metabolic end-product.^{13,18,19,22–24}

While many members of the order grow lithotrophically, most known species exhibit modes of either strict heterotrophy or mixotrophy. Most members of the genera *Sulfolobus* and *Metallosphaera* are capable of utilizing protein-rich substrates, such as yeast extract or tryptone, under aerobic conditions. Furthermore, several species, such as *Sulfolobus solfataricus*, *Sulfolobus shibitae*,²⁵ and *Sulfolobus islandicus*,²⁶ use a wide variety of sugars in catabolic metabolism. In addition, members of the order, particularly in the genera *Metallosphaera* and *Acidianus*, are capable of oxidizing metal sulfides: a trait that is particularly useful for bioleaching of base, precious and strategic metals from mineral ores.^{27–30} Finally, some members of the genus *Acidianus* are capable of using ferric iron as an electron acceptor under anaerobic conditions.^{21,31}

Carbon Dioxide Fixation

Interestingly, the natural habitats of many Sulfolobales (solfatara/calderas) are limited or devoid of complex carbon sources, necessitating the process of autotrophy. The ability of organisms to fix carbon dioxide from the atmosphere is considered by many to explain the early formation of the multi-carbon molecules required to fuel life, explaining their retention in species across all three domains of life.^{32,33} As it stands, 6 major routes exist for the fixation of carbon dioxide: the Calvin-Benson cycle (present in most plants), the reductive citric acid cycle (green sulfur bacteria), the reductive acetyl-CoA cycle (acetogens/methanogens), the hydroxypropionate bi-cycle (*Chloroflexus*), and the 3-hydroxypropionate/4-hydroxybutyrate (3-HP/4-HB) or dicarboxylate/4-hydroxybutyrate pathways (both from the Crenarchaeota).³⁴ The habitats from which the Sulfolobales were isolated, unlike many

other organisms, are characterized by copious oxygen and inorganic electron donors.¹² While the reactions driving this cycle are some of the most energy-demanding for autotrophic carbon assimilation, their advantage may lie in their relative insensitivity to oxygen, avoidance of side-reactions, direct utilization of bicarbonate, and thermal stability.^{34,35}

Briefly, the 3-HP/4-HB cycle has two major products that enter into cellular metabolism. The first portion involves the addition of two bicarbonate molecules to acetyl-CoA to form succinyl-CoA, which is subsequently reduced in the second half of the cycle to 4-hydroxybutyrate and eventually dissociated to two molecules of acetyl-CoA (Figure 1).³⁶ The key enzyme in the cycle is a biotinylated acetyl-CoA/propionyl-CoA carboxylase, that is, bifunctional and efficient in substrate turnover.³⁷ Metabolic analysis of the cycle has revealed that the major product of the cycle is not acetyl-CoA (as originally hypothesized), but rather succinyl-CoA (roughly two-thirds of the carbon flux), yielding malate and oxaloacetate in subsequent oxidation reactions.³⁸ This requires a turn and a half of the cycle to maintain acetyl-CoA levels and generate succinyl-CoA.

From an application-oriented point-of-view, it may be possible to use this pathway to sustainably produce high-value specialty chemicals, such as 3-hydroxypropionate (3-HP) or succinate. The former is used industrially in polymer production and the latter is used to produce solvents and polymers.⁴⁶ For this reason, several attempts have been made to utilize these genes in metabolically engineered hosts. For instance, the first three enzymes have been expressed in *P. furiosus* to introduce a temperature-shift-responsive metabolic mode for the production of 3-HP.⁴⁷ Further work with this metabolically engineered strain has demonstrated that the addition of a biotin protein ligase can improve 3-HP titers more than eight-fold.⁴⁸ This dramatic improvement is likely due to the presence of a biotinylated subunit on the key acetyl-CoA carboxylase enzyme from the cycle.³⁶ Thus, this well-studied pathway has opportunities to be utilized and improved upon.

Sulfur Utilization

In contrast to carbon metabolism, sulfur metabolic pathway discovery is hampered by the tendency of elemental sulfur and its derivatives to react nonenzymatically, masking the true identity of an enzyme's substrate or products.^{49,50} Because S^0 is sparingly soluble in water under standard conditions, the true substrate for microbial growth on sulfur is likely

soluble polysulfides and polythionates, introduced by nonenzymatic reactions.^{51–53} However, these solubilizing reactions only occur at near-neutral pH, since under acidic conditions the equilibrium strongly favors elemental sulfur.⁵⁰ In order to overcome these solubility issues, it has been proposed that acidophiles may physically associate with sulfur particles or that at elevated temperatures sulfur becomes sufficiently soluble to support growth.⁵³ Regardless, these organisms have a suite of proteins capable of manipulating the initial elemental sulfur from the environment, as well as many of its derivatives.

The sulfur oxygenase reductase (SOR), first identified in a member of the Sulfolobales, *Acidianus ambivalens*,⁵⁴ appears to be key to acidophilic sulfur oxidation. This intracellular enzyme is active on elemental sulfur, indicating transport of elemental sulfur or one of its derivatives to the cytoplasm by some still unidentified mechanism. SOR appears to be limited to the Sulfolobales and a few extremophilic bacteria;⁵⁵ this makes sense given that other sulfur lithotrophs grow closer to neutral pH where more soluble sulfur species are abundant. SOR acts on elemental sulfur by disproportionating it equally between oxidized (sulfite; SO_3^{2-}) and reduced (hydrogen sulfide; H_2S) products. Further, the production of thiosulfate observed in early studies of the SOR⁵⁴ is now believed to be the result of a nonenzymatic reaction.⁵⁶ SOR requires oxygen to function but uses no additional co-factors, suggesting its ability to conserve cellular energy. Instead, it 'activates' long, unwieldy hydrophobic sulfur chains into smaller intermediates that can be used by other enzymes to generate energy.

Acidianus ambivalens has served as the model organism for sulfur metabolism studies in the Sulfolobales, since measurements of sulfur-active enzymes from cell extracts were used to construct a conceptual model of its sulfur oxidation pathways.⁵⁷ While the SOR enzyme has been the most thoroughly characterized enzyme with respect to sulfur metabolism,^{54,55,58} studies of enzymes purified or detected from *A. ambivalens* cell extracts provide some insights into the complete oxidation pathway. The *A. ambivalens* genome remains unsequenced, so many of its enzymes are identifiable only by their activity in cell extracts. This presents a challenge for systems biology-based efforts to understand the details of sulfur oxidation in other Sulfolobales, or even how this process contributes to *A. ambivalens* energetics and metabolism. Regardless, there appears to be two parallel processes by which




FIGURE 1 | 3-Hydroxypropionate/4-hydroxybutyrate (3-HP/4-HB) cycle from *Metallosphaera sedula*. The cycle consists of two major portions: carbon incorporation (via bicarbonate) occurs in the first half (blue) of the cycle and is followed by subsequent reduction and reformation of two acetyl-CoA molecules in the second half (red). Enzymes listed and their references: acetyl-coA carboxylase^{37,39} (ACC), acetoacetyl-CoA β -ketothiolase³⁶ (ACCT), acryloyl-CoA reductase⁴⁰ (ACR), crotonyl-CoA hydratase⁴¹ (CCH), 4-hydroxybutyrate-CoA dehydratase³⁶ (HBCD), 4-hydroxybutyrate-CoA synthase⁴² (HBCS), 3-hydroxypropionate-CoA dehydratase⁴⁰ (HPCD), 3-hydroxypropionate-CoA synthase⁴³ (HPCS), methylmalonyl-CoA epimerase⁴⁴ (MCE), methylmalonyl-CoA mutase⁴⁴ (MCM), malonyl-CoA/succinyl-CoA reductase⁴⁵ (MCR), malonate semialdehyde reductase⁴⁵ (MSR), and succinate semialdehyde reductase⁴⁵ (SSR).

A. ambivalens (and presumably the other sulfur-oxidizing members of the Sulfolobales) gain energy while oxidizing elemental sulfur to sulfate (SO_4^{2-}). One pathway uses the membrane-associated oxidoreductases, TQO and SAOR, to reduce an electron carrier (such as quinone),⁵⁹ thereby generating proton motive force via the terminal oxidase,^{60,61} while the other pathway produces one high-energy phosphate bond (ADP from AMP) by direct substrate level phosphorylation via APSR

and APAT, generating sulfate in the process⁵⁷ (Figure 2). While the sulfur metabolism has been examined in bioleaching applications (see next section), the ability of *S. metallicus* to remove toxic H_2S from high-temperature gas streams represents a potentially important technological use of sulfur oxidation.⁶²

A. ambivalens has also served as the model system for anaerobic sulfur reduction among the Sulfolobales.⁶³ The enzyme pathway for sulfur reduction




FIGURE 2 | Chemolithotrophic pathways in the Sulfolobales. The first half of the figure (blue) shows the hypothetical pathways for sulfur utilization in the Sulfolobales, including both oxidizing and reducing pathways, beginning with elemental sulfur. Sulfur reducing complexes: hydrogenase (Hyd), sulfur reductase (SR).⁶³ Sulfur oxidizing enzymes: sulfur oxygenase reductase (SOR),⁵⁴ thiosulfate:quinone oxidoreductase (TQO),⁵⁹ Sulfite:acceptor oxidoreductase (SAOR), adenylylsulfate reductase (APSR), adenylylsulfate:phosphate adenylyltransferase (APAT).⁵⁷ The second panel shows a hypothetical pathway for the oxidation of ferrous iron using several *fox* stimulon proteins as well as some iron-responsive respiratory proteins. Ferrous-oxidation (Fox), rusticyanin (Rus), cystathionine- β -synthase containing protein subunits A and B (CbsAB), sulfur oxidation (Sox), NADH dehydrogenase (NAD).^{29,64,65}

in *A. ambivalens* appears to be much simpler than for oxidation, possibly involving only two closely associated membrane complexes. A membrane hydrogenase passes electrons (via a quinone molecule) from H_2 to a sulfur reductase, where they are used to reduce elemental sulfur to H_2S .⁶³ The cycling of quinones between the two enzymes—forming a ‘redox loop’ similar to the one used in *Escherichia coli* during growth on nitrate⁶⁶—is likely the way protons are transported across the membrane, coupling sulfur reduction to energy conservation (Figure 2).

Metal Oxidation

Along with interest in sulfur metabolism, some of the earliest work in determining the mechanism of metal oxidation in the Sulfolobales (and acidophiles, in general) involved the spectroscopic identification of unique cytochromes from iron-oxidizing cultures.⁶⁷ This original research led to the intensive study and eventual development of a model in the mesoacidophile *Acidithiobacillus ferrooxidans*, involving the shuttling of electrons from the outer membrane of the cell to the inner membrane, driving a terminal oxidase to maintain pH homeostasis and the production of reducing power for intracellular metabolic needs.⁶⁸ Not surprisingly, the spectroscopic data from *S. metallicus* demonstrated early on that key differences exist between bacterial and archaeal metal oxidation, particularly in the presence of cytochromes and the roles of various protein

complexes in transporting electrons.^{67,69} However, some of the same systems-based approaches were utilized to detect the transcriptomic response of known iron-oxidizers, including *S. metallicus*,⁶⁴ *M. yellowstonensis*⁷⁰ and *M. sedula*,⁶⁵ in the presence of iron. Interestingly, these experiments suggest the importance of merging several systems biology techniques in order to ascertain new pathways and information. While all three species contain the *fox* stimulon (an assortment of ferrous-responsive genes A–J) and key related genes (such as rusticyanin and the cystathionine- β -synthase subunits A and B), their regulation varies dramatically among the species with both constitutive and inducible expression observed during iron supplemented growth.^{64,65,70} Yet, the merger of this transcriptomic data and genomics analysis yielded a hypothesized pathway for metal biooxidation in these organisms (Figure 2), which relies on a cytochrome b (as opposed to a cytochrome c), bifurcating rusticyanin(s), and two terminal complexes: an NADH dehydrogenase (generating reducing power) and a putative cytochrome c oxidase (driving pH homeostasis). Although similarities exist between the two systems in *A. ferrooxidans* and *Metallosphaera/Sulfolobus spp.*, distinctive co-factors and apparent differences in organization suggest that these systems are evolutionarily divergent modes of biooxidation.⁷¹

These differences, as well as the major phenotypic differences between these two classes of metal mobilizers, relates to their use in metal bioleaching applications. For example, many mesophilic organisms

are ill-suited to bioleaching of highly gangue (i.e., high sulfur content) ores due to the extremely exothermic nature of sulfur oxidation chemistry. The build-up of heat can be problematic in large heap operations that rely on mesophiles alone.^{72,73} This physiological trait cannot be undervalued given that the removal of elemental sulfur can improve cyanidation (a form of chemically-driven solubilization), which is commonly used in gold mining. Furthermore, extreme thermophiles appear to have some niche advantages over mesophiles for bioleaching of several specific types of copper ores, including the enhanced dissolution of copper from recalcitrant primary ores (such as chalcopyrite),^{74–76} selective mobilization of copper over molybdenum in copper-bearing molybdenite,^{77,78} and the unassisted mineralization of arsenic in the form of arsenate from enargite ores.⁷⁹ Bioleaching operations targeting copper have increased dramatically and currently account for more than 15% of the global output.⁸⁰ Thus organisms that present an inherent propensity for copper solubilisation such as *A. brierleyi*, *S. metallicus*, or *M. sedula* deserve more investigation for their potential industrial application.

PYROCOCCUS FURIOSUS

Pyrococcus furiosus, the type strain of the genus, was first isolated in 1986 from a hydrothermal vent off of the coast of Vulcano Island (Italy) and has been one of the most studied hyperthermophiles to date, due to its intriguing phenotypical characteristics.⁸¹ Exhibiting optimal growth at 100°C and a pH near 7, it was the second genus, after the autotrophic sulfur-oxidizing *Pyrodictium*, capable of growth at temperatures at or above 100°C.⁸¹ As a heterotrophic organism, *P. furiosus* is capable of utilizing hexose oligosaccharides such as cellobiose and laminarin,⁸² chitin,^{83,84} and peptides.⁸⁵ Efforts over the past three decades have elucidated many unique features of this organism, including various novelties in metabolic pathways, regulatory mechanisms and proteins and enzymes.

Central Glycolytic Metabolism

P. furiosus grows well on disaccharides (maltose and cellobiose) and glucans (laminarin and starch), but not on monosaccharides nor pentoses.⁸⁶ The reasons for this anomaly are unknown but monosaccharides may not be available externally in these high-temperature environments as they are susceptible to the Maillard reactions, in which sugars react with

available amino acids to form glycosylamines; this problem is especially exacerbated in peptide-rich media. Disaccharide and polysaccharide transport may also be more efficient energetically.

P. furiosus derives no net substrate level phosphorylation from glucose to pyruvate conversion, unlike the traditional Embden-Meyerhof (EM) pathway that provides two ATP per glucose and the Entner-Doudoroff (ED) pathway which yields one ATP per glucose (see Figure 3). Thus, the only net substrate level phosphorylation gains are a result of ATP-forming hydrolysis of acetyl-CoA, produced from pyruvate via pyruvate oxidoreductase and acetyl-CoA synthetase.⁸⁷ *P. furiosus* contains a nontraditional variation of EM glycolysis, in which glucokinase and phosphofructokinase utilize ADP as the phosphoryl group donor, generating AMP.⁸⁶ In the early 1990s, these were the first reported ADP-dependent kinases.^{86,88}

The absence of an energy-conserving step in the glycolytic pathway is due to the absence of a 1,3-biphosphoglycerate intermediate, which is found in both the EM and ED pathways. As shown in Figure 3, this direct conversion from glyceraldehyde-3-phosphate (GAP) to 3-phosphoglycerate (3PG) does result in production of a reducing equivalent in the form of reduced ferredoxin, but does not result in substrate level phosphorylation. The phosphate group is released via hydrolysis without capture of this high-energy bond. The enzyme responsible for the conversion of GAP to 3PG, GAP ferredoxin oxidoreductase (GAPOR), is unusual in that it requires tungsten, an element rarely found in biology.⁸⁹ The absence of other, more traditional glycolytic enzymes makes GAPOR's function critical to sugar utilization. Thus, tungsten levels have a significant impact on the growth of *P. furiosus* in the presence of maltose.⁸⁹

Fermentation Pathways

P. furiosus produces reduced ferredoxin through central glycolysis and, as an obligate anaerobe, must have a route to dispose of any excess reducing power. Two primary routes exist for this purpose depending on hydrogen partial pressures, the availability of elemental sulfur and nitrogen, and other regulatory factors. The primary route of regenerating oxidized ferredoxin is through a membrane-bound hydrogenase (MBH) that produces an ion gradient that allows ATP production via ATP synthase.⁹⁰ The hydrogenase is thought to exchange the proton gradient generated by hydrogen production for a Na⁺ gradient and this is utilized by a Na⁺-dependent ATP synthase.⁹¹ This energy-conserving hydrogenase




FIGURE 3 | Comparison of traditional Embden-Meyerhof-Parnas pathway with the modified pathway in the archaeon *P. furiosus*. Included are three fermentative pathways which utilize the reduced ferredoxin produced via glycolysis. Enzyme abbreviations: hydrogenase (hyd), ferredoxin: NADP oxidoreductase (FNOR), glutamate deaminase (GD), alanine aminotransferase (AT), and NADP:sulfur oxidoreductase (NSOR).

therefore constitutes a single-step electron transport chain, and has been proposed as an evolutionary precursor to the complicated, multi-step electron transport chains that are common in present day microbes.⁸⁷ While the exact mechanism coupling proton transfer and hydrogen production is unknown, it is estimated that 0.3–0.4 molecules of ATP are produced per two electrons transferred.⁸⁷ Thus approximately 1.2 moles of ATP are produced

for every mole of glucose converted to acetate via glycolytically produced reducing equivalents. Given the low energy production resulting from glycolysis, this fermentative process is particularly critical.⁸⁷

When elemental sulfur is present, *P. furiosus* produces hydrogen sulfide rather than hydrogen gas.⁹² As with H₂ production, a proton gradient is formed by a membrane-bound oxidoreductase (MBX).⁹³ MBX is highly homologous to the MBH

and is thought to oxidize ferredoxin.⁹³ However, it is not known if MBX reduces sulfur directly or generates NADPH that is then used by a cytoplasmic NADPH- and CoA-dependent enzyme.⁹³ The reason for the preference for sulfur over proton reduction is not clear but it is strong since the switch from H₂ to H₂S production begins only minutes after the addition of sulfur to a growing culture.⁹⁴ The shift is mediated by SurR, a redox-responsive transcriptional regulator that has been well characterized.^{95,96}

Another method for disposing of reductant during fermentation involves the transformation of pyruvate to alanine with the addition of available nitrogen.⁹⁷ This results in a major energetic penalty, however, as the pyruvate is not used to produce acetyl-CoA, which is responsible for the majority of ATP production. Thus, alanine pathway is only utilized when sulfur is absent and the hydrogen partial pressures are high.⁹⁷

Applied Biocatalysis and Metabolic Engineering

Prior to detailed knowledge of the *P. furiosus* genome and development of genetic manipulation methodology, early work focused on characterizing its novel enzymes, with an eye toward industrial applications. While the DNA polymerase from *Thermus aquaticus* (Taq) is the most widely known and utilized thermostable polymerase in PCR reactions, the *P. furiosus* DNA polymerase is considerably more thermostable and of higher fidelity, yet two- to threefold lower processivity. Owing to its 3' to 5' exonuclease proofreading activity, the polymerase exhibits a 10-fold reduction in error rate compared to the Taq polymerase.⁹⁸ Additionally, the NADP(H)-dependent hydrogenase (SH1) from *P. furiosus* is extremely thermostable and has a temperature optimum of 95°C.⁹⁰ It has been utilized in a renewable H₂ production *in vitro* system in which sugars are completely oxidized to CO₂ and H₂. A combination of pure enzymes comprising the pentose phosphate pathway (PPP) were used to convert sugars to CO₂ and the NADPH that is then produced is oxidized and H₂ is produced by SH1.⁹⁰ Many other enzymes of interest from *P. furiosus* have been purified and characterized,⁹⁹ and include carbohydrate hydrolyzing enzymes (e.g., α -amylase,¹⁰⁰ amylopullulanase,¹⁰⁰ endoglucanase,¹⁰¹ and β -glucosidase,¹⁰² and chitinase⁸³) and proteases.^{103,104}

P. furiosus now has a facile genetic system which has led to efforts directed at metabolic engineering.¹⁰⁵ Earlier work on a related extreme thermophile, *Thermococcus kodakarensis*, a member of

the same order as *P. furiosus*, the Thermococcales, paved the way for the *P. furiosus* genetic tools.¹⁰⁶ For *P. furiosus*, its high growth temperature and tolerance to cold shock opens up its use for hosting metabolic pathways from much less extreme thermophiles.^{107–109} In fact, a novel temperature-shift strategy has been demonstrated that minimizes *P. furiosus* metabolism at sub-optima temperatures to direct energy to heterologous product formation.¹¹⁰ As mentioned above, *P. furiosus* produces soluble hydrogenases, which can regenerate reducing equivalents from hydrogen gas.⁴⁷ These hydrogenases could allow metabolically engineered *P. furiosus* to use electrons from H₂ to produce highly reduced chemical products.⁴⁷ The insertion of three genes from the *M. sedula* 3-HP/4-HB carbon fixation cycle into *P. furiosus* demonstrated production of 3-HP utilizing sugars and sequestering carbon dioxide for a portion of the molecule.⁴⁷ Another heterologous pathway expressed in *P. furiosus* utilized genes from three thermophilic organisms with optimal temperatures ranging from 65 to 75°C for the production of *n*-butanol.¹¹¹ With this alcohol pathway, significant diversion to ethanol was shown due to promiscuity of the aldehyde dehydrogenase enzymes.¹¹¹ The use of less thermophilic enzymes in heterologous biosynthetic pathways has provided insights into *P. furiosus* native metabolism at lower temperatures. For example, at 70–80°C, acetoin is produced as a major metabolic product. Along these lines, it was shown that the removal of acetolactate synthase in *P. furiosus* generates small amounts of ethanol as a metabolic end product, as pyruvate was directed toward acetate and eventually ethanol rather than acetoin.¹⁰⁹ In addition to its native abilities to utilize sugars, *P. furiosus* was engineered with a 16 gene cluster to oxidize carbon monoxide to carbon dioxide, producing H₂ and energy in the process.¹⁰⁷ Overall, the ability to engineer utilization of unique energy sources, manipulate temperatures to optimize enzyme activities, and insert genes from a variety of organisms with different optimal growth temperature provide tools not typically available in model mesophilic hosts.

CALDICELLULOSIRUPTOR SPP.

Caldicellulosiruptor is a bacterial genus containing the most thermophilic, cellulolytic microorganisms known to date. Isolated worldwide and having optimal growth temperatures between 70 and 78°C, these Gram-positive, asporogenic, obligate anaerobes have the ability to degrade untreated lignocellulosic

biomass, a highly sought after phenotype for consolidated bioprocessing of fuels and chemicals. Many well-studied cellulolytic microbes are known to either secrete individual enzymes or large cellulolytic enzyme complexes (e.g., the cellulosome¹¹²) into their environment. In contrast, *Caldicellulosiruptor* species instead use an array of multi-modular enzymes to breakdown plant biomass.^{113–118} These carbohydrate-active enzymes (CAZymes) are composed of both catalytic [e.g., glycoside hydrolases (GH)] and noncatalytic [e.g., carbohydrate binding module (CBM)] domains. All *Caldicellulosiruptor* species are able to utilize fructose, galactose, glucose, xylose, and pectin via a classical EMP pathway.^{119–121} However, arabinose, rhamnose, and fucose utilization, is not conserved throughout all species.^{119,121–124} Some of the sugars, such as xylose and arabinose, are broken down via the nonoxidative PPP and then piped into the EMP pathway as intermediates.¹²⁵ Although *Caldicellulosiruptor* species lack the oxidative PPP, which generally is responsible for NADPH production, they are still capable of generating NADPH; the exact enzymatic mechanism for this process is currently unknown.¹²¹ Members of the *Caldicellulosiruptor* genus also contain an incomplete tricarboxylic acid (TCA) cycle, consisting of a reductive branch leading to fumarate and an oxidative branch producing succinyl-CoA.

Carbohydrate Utilization

Caldicellulosiruptor spp. produce many highly versatile and efficient multi-modular carbohydrate-degrading enzymes, made up of combinations of GH and CBM domains. For example, CelA, a lignocellulosic CAZyme, is composed of five carbohydrate-specific domains: GH9-CBM3-CBM3-CBM3-GH48 connected by linker regions.^{126–128} These different segments allow proteins to have multiple functions: simultaneously binding to its substrate (via the CBM3s), as well as cleavage of specific bonds (via the GH9 and GH48 domains); these GH9 and GH48 domains are capable of endo- and exoglucanase activity. CelA is present in only the most cellulolytic species of the genus and its two GH domains provide these species with a unique ‘drilling’ mode of action during biomass deconstruction.¹²⁶ While CBMs allow the CAZyme complex to adhere to the biomass, surface layer homology (SLH) domains are also found in the *Caldicellulosiruptor* multi-modular scheme.^{129,130} Instead of being freely transported out of the cell, CAZymes with SLH domains are tethered to the cell’s S-layer. As such, the enzymes can breakdown and bind, if they contain CBMs, substrates in

close proximity to the microbe, providing better access to available sugar moieties. It also has been recently found that *Caldicellulosiruptor* species have a novel method of attaching themselves to crystalline cellulose. Structurally unique proteins, called tāpirins, are expressed on the cell surface, and contain a binding domain specific to insoluble cellulose.¹³¹ Present in every member of the genus and highly expressed, tāpirins are thought to play an important role in how plant matter is deconstructed by microbes this genus.

Both methods of attachment, along with the large inventory of glycolytic enzymes, give this genus its impressive ability to degrade a wide variety of lignocellulosic substrates.¹¹² *Caldicellulosiruptor* species are capable of breaking down cellulose and hemicellulose (hexoses and pentoses), both as simple monosaccharides and complex biomasses.^{114–116,122,132–136} Unlike many cellulolytic organisms, they do not exhibit carbon catabolite repression, a process by which certain sugars are preferentially metabolized, while excluding the usage of others.^{112,117,120} This is especially advantageous in an industrial process involving lignocellulose conversion to fuels and chemicals, as these microbes can utilize multiple sugars simultaneously, with numerous points of entry to central carbon metabolism (see Figure 4). Although *Caldicellulosiruptor saccharolyticus* was shown to grow well on a variety of sugars (arabinose, fructose, galactose, glucose, mannose, and xylose) simultaneously, the extent of which each monosaccharide was digested varied, with fructose being the most utilized.¹¹⁷ In the absence of an apparent carbon utilization regulatory system, variation in sugar utilization among *Caldicellulosiruptor* species is likely due to presence or absence of certain metabolic pathways, for example, the oxidative PPP, and/or essential transporters; the latter has only recently been better understood for a few *Caldicellulosiruptor* species with transcriptomics analysis of growth on substrates, such as simple sugars, crystalline cellulose (Avicel), and complex biomasses like switchgrass.^{113,114,117,121,137}

One option to improve degradation of lignocellulose is to increase the CAZyme inventory of a microbe. While generally highly conserved in the *Caldicellulosiruptor* genus, the SLH domain xylanase from *Caldicellulosiruptor kronotskyensis*, Calkro_0402, is not present in *C. bescii* and, thus, was inserted into the genome to improve its ability to utilize xylan.¹³⁰ The manipulated strain successfully expressed the protein on the S-layer of the cell and improved xylan utilization significantly by doubling xylose release into the supernatant from oat spelt xylan. Growth on washed and unwashed birch xylan was improved, while dilute acid-pretreated switchgrass solubilization remained




FIGURE 4 | Conserved metabolic pathways in all *Caldicellulosiruptor* species. This includes sugar uptake, glycolytic, and fermentative pathways. The figure includes only the major steps, or start and end products. P is the abbreviation for phosphate, NADH for reduced nicotinamide adenine dinucleotide, ATP for adenosine triphosphate, Fd^{red} for reduced ferredoxin, and GDP for guanosine diphosphate.

unaffected; this indicated that there are still other hurdles to lignocellulosic degradation that must be overcome. However, on substrates with high xylan content, the engineered strain showed improved solubilisation, possibly due to increased substrate attachment.

Fermentation

The major fermentation products of the *Caldicellulosiruptor* genus are hydrogen, carbon dioxide, and acetate.^{122,138,139} Lactate production has also been measured, but only a trace amount of ethanol has been detected in wild-type cultures.^{119,123,124,140} By far, the most studied product of all of these is molecular hydrogen, especially with *C. saccharolyticus*.^{120,121,141–154} H_2 generation is completed via hydrogenases utilizing reducing equivalents (Fd^{red} and NADH) from central carbon metabolism (see Figure 4). *C. saccharolyticus*, along with several other extreme thermophiles, is

considered to be an ideal ‘biohydrogen factory’, as reported yields are close to the so-called ‘Thauer limit’ of 4 moles H_2 per mol glucose.^{120,122,155} Decreased H_2 production results in the accumulation of NADH and Fd^{red} , while increased H_2 can instead push metabolic flux toward lactate production. At high levels of molecular hydrogen, product inhibition occurs via increased dissolved H_2 levels.¹⁵⁶ NADH and Fd^{red} are simultaneously oxidized by a bifurcating [FeFe]-hydrogenase, which uses both electron donors at the same time,¹⁵⁷ while Fd^{red} is also oxidized by a membrane-bound [NiFe]-hydrogenase that is related to that found in *P. furiosus*.⁹¹

With the recent development of a genetic engineering system in *Caldicellulosiruptor bescii*, based on auxotrophic selection-targeted manipulations of the *Caldicellulosiruptor* genome and consequently metabolism are now possible.^{158–161} The first directed demonstration of these methods actually involved the deletion of the single lactate dehydrogenase

(*ldh*—Athe_1918) present in *C. bescii* to halt lactate production.¹⁶² While the wild type and parent strains produced less hydrogen than the well-studied *C. saccharolyticus* (1.8 and 1.7 versus 2.5 mol H₂/mol of glucose, respectively), the *ldh* knockout produced significantly more H₂ on switchgrass, closer to the theoretical goal (3.4 mol H₂/mol of glucose). As lactate formation ceased, acetate production increased by 38–40% over the wild type and parent strains.

More recently, ethanol production was demonstrated in *C. bescii* through the addition of an NADH-dependent alcohol dehydrogenase gene from *Clostridium thermocellum* (*adhE*—Cthe_0423) into the strain lacking lactate formation¹⁶³; *C. bescii* does not possess a native alcohol or acetaldehyde dehydrogenase, and thus a representative gene was recruited from another thermophilic Firmicute. Owing to the lower thermostability of the *C. thermocellum* protein, growth of the engineered strain was done at a maximum of 65°C. Strain growth on cellobiose, Avicel and switchgrass, and resulted in 14.8 mM, 14 mM, and 12.8 mM ethanol, respectively. Acetate production was also lowered, ranging from ~4 to 5 mM compared to the wild type (~6 mM) and parent (~8–9 mM) on all tested substrates. Another attempt at ethanol production was completed by individually inserting two bi-functional alcohol dehydrogenase genes from *Thermoanaerobacter pseudethanolicus* 39E, *adhB* (Teth39_0218) and *adhE* (Teth39_0206), into the *C. bescii* lactase dehydrogenase knockout.¹⁶⁴ Growing the modified strain at 75°C with cellobiose, ethanol was produced at reported levels of 1.4 mM and 2.9 mM, acetate at 15.5 mM and 14.1 mM, and H₂ at 23.2 mM and 22.5 mM for the AdhB and AdhE knock-in strains, respectively; similar ethanol levels were also measured on Avicel and switchgrass.

With a genetic system now in place, gene ‘knockouts’ in *C. bescii* can be strategies to understand *Caldicellulosiruptor* metabolism and physiology. For instance, an uncharacterized [Ni-Fe] hydrogenase maturation gene cluster (*hypABFCDE*—Athe_1088–Athe_1093) was deleted from the aforementioned, modified ethanol-producing *C. bescii* strain containing *adhE* from *C. thermocellum*.¹⁶⁵ The resulting strain produced 20% less H₂ than its parent, yet its H₂ yield per mol of cellobiose increased 63% (3.58 versus 2.19 mol H₂/mol cellobiose). Fermentation patterns on Avicel, cellobiose, and switchgrass showed that the engineered strain also produced acetate (1.6–5.7 mM—34% less than parent) and ethanol (1.9–2.7 mM—73% less than parent). Additionally, the knockout

had reduced growth and a longer lag phase, which could result from the deleted gene acting as an ATP-generating protein pump. Other genetic manipulations with *C. bescii* include the deletion of *CelA* (Athe_1867),¹²⁸ a predicted pectin lysase, and a putative AraC family transcriptional regulator genes (*pecABCR*—Athe_1853–1856).¹⁶⁶ Continued efforts with the newly established genetics tools in *C. bescii*, and eventually other *Caldicellulosiruptor* species, will help reveal the basis for its ability to grow on lignocellulosic substrates.

THERMOTOGA SPP.

The bacterial genus *Thermotoga* contains nine named species that are obligate anaerobes capable of growth at optimal temperatures between 65 and 80°C, mostly isolated from submarine geothermal features.^{167–169} These rod-shaped, Gram-negative, eubacteria were originally identifiable by their distinctive ‘toga’-like outer sheath and absence of an outer membrane. Beyond their unique appearance, the species in the genus *Thermotoga* share a remarkably large number of homologs (roughly 24% of the genome) with sequenced archaea.¹⁷⁰ This curious result has led some research into the evolutionary divergence/convergence of this bacterial lineage, suggesting the genomic features that may be critical in defining thermophily, such as the discovery of genes associated with biosynthesis of di-myoinositol-phosphate, which may serve as a critical thermoprotectant compatible solute.¹⁷¹ Further phylogenetic analysis has even suggested that mesophily may have developed from thermophily (within the order Thermotogales), given the ancestral sequence reconstruction of more thermally stable myo-inositol-phosphate synthase (MIPS)¹⁷² and emergence of ‘mesotoga’ species.¹⁷³ Also of interest is the presence of a system for catabolizing myo-inositol that provides utilization of compatible solutes but cannot provide a complete source for carbon utilization.¹⁷⁴ Within the genus, *Thermotoga maritima* has served as a model species for studying evolution, biomass deconstruction, and biohydrogen production.¹⁷⁵

Carbohydrate Utilization

All *Thermotoga* species are chemoheterotrophs, although the range of substrate usage varies and includes numerous pentoses, hexoses, disaccharides, and polysaccharides, as well as yeast extract, acetate, methanol, and pectin.^{168,169,176–180} This ability to utilize a broad array of carbohydrates appears to be supported by bioinformatics and transcriptomics

suggesting a substrate-specific regulation and function of large numbers of ABC-transporters,¹⁸¹ as well as many α - and β -GH.¹¹⁸ Intriguingly, *T. maritima* grows faster on complex carbohydrates than on monosaccharides, suggesting an adaptation to the breakdown of biomass in their natural environments.¹⁸² The metabolism of carbohydrates by these organisms results not only in the formation of some typical fermentation products, such as acetate, carbon dioxide, and lactate, but also the generation of molecular hydrogen, and small amounts of ethanol, butanol, and butyrate.¹⁸³ It is worth noting that *Thermotoga* species utilize the traditional EMP and ED pathways^{184,185} for carbon utilization. However, they also contain, in some cases, unique enzymes that are adapted to optimizing the use of reducing power and energy generated from biomass deconstruction for the synthesis of fermentation products.

The initial genome annotation of *T. maritima* suggested a prevalence of mono- and polysaccharide utilization proteins (as much as 7% of identified genes).¹⁷⁰ In contrast to organisms that produce large complexes for carbohydrate degradation (i.e., cellulosomes), *T. maritima* utilizes a broad array of both extra- and intra-cellular GH, which have been detailed in previous reviews.¹¹⁸ More recent examination of the pan genome, as well as transcriptomic data, suggests that *Thermotoga* species vary with respect to specific ABC sugar transporters and GH.¹⁸⁶ Overall, the preponderance of thermally stable, polysaccharide-degrading enzymes makes members of the genus and their enzymes intriguing candidates for the deconstruction of complex carbohydrates in industrial applications.¹⁸⁷ However, one of the limiting factors is the absence of any apparent capacity for growth on crystalline cellulose, suggesting a lack of cellulolytic enzymes in *Thermotoga* species.¹⁸⁸ In fact, to address this issue, efforts were directed at the ectopic expression of cellulases from *C. saccharolyticus* fused with *T. maritima* signal peptides. The resulting plasmids were used for *Thermotoga* sp. strain RQ2 transformations, where enhanced exoglucanase activity was observed, but eventually was lost due to poor plasmid maintenance.¹⁸⁹ However, a stable genetic system for *T. maritima* and *T. sp.* RQ7 was recently reported, based around a cryptic plasmid isolated from the latter.¹⁹⁰

Fermentation

Of the major fermentation products from *Thermotoga* spp., H₂ production is particularly interesting from a biotechnological perspective. High yields

(3.8 mol H₂/mol glucose) reported by *Thermotoga neapolitana* under anaerobic and microaerobic growth conditions^{191–193} approach the Thauer limit.¹⁵⁵ The production of H₂ is most efficient when the balance of fermentation products is skewed toward acetate production as compared to lactate production, given that the enzymes identified in acetate production, phosphate acetyl-transferase and acetate kinase, avoid the re-oxidation of NADH and instead produce Fd^{red} and ATP, respectively. In contrast, the production of lactate is driven by a lactate dehydrogenase that uses reducing equivalents (NADH) generated in the glycolytic process. Another possible key to efficient hydrogen production in these organisms, as in the *Caldicellulosiruptor*, is the coupling of Fd^{red} and NADH oxidation by a bifurcating, [FeFe] hydrogenase,¹⁹⁴ in which Fd^{red} likely drives the less favorable oxidation of NADH and improves the overall thermodynamics for producing hydrogen. This enzyme complex, first identified in *T. maritima*, appears to have a homolog in *T. neapolitana*, which has the highest reported H₂ yields within the genus. There is also evidence that the build-up of molecular hydrogen and a possible inhibition mechanism can be alleviated through the co-culturing of *T. maritima* with *Methanococcus jannaschii*; the latter oxidizes H₂ and generates methane.¹⁹⁵ This results in significant upregulation of CAZymes and growth-phase enzymes, as well as denser cultures¹⁹⁶ (see Figure 5).

Besides molecular hydrogen, several species have been reported to produce ethanol as a fermentation product.^{177,197} This result was not expected given the lack of detectable pyruvate decarboxylase activity. However, more recent work has identified the presence of both an alcohol dehydrogenase (from *Thermotoga hypogea*)¹⁹⁸ and a bi-functional




FIGURE 5 | Co-culture of *Thermotoga maritima* yellow/green rods and *Methanocaldococcus jannaschii* (red cocci)¹⁹⁶—permission pending.

pyruvate ferredoxin oxidoreductase-pyruvate decarboxylase enzyme.¹⁹⁹ Additionally, butyrate (an odiferous organic used primarily as a perfume or food additive) production has been linked with hydrogen biosynthesis in studies involving *T. neapolitana*²⁰⁰; however, the mechanism of butyrate synthesis is still unknown in these organisms.

THERMUS SPP.

The genus *Thermus* was among the first bacteria to be studied with respect to thermophily,²⁰¹ and the DNA polymerase from *Thermus aquaticus*² was used in early efforts with the PCR. While not all *Thermus* species (at least 11 have been named and characterized) grow optimally at 70°C and above (extremely thermophilic), there are some that meet this thermal threshold. *Thermus* species are typically nonmotile, nonsporulating, and are not naturally capable of fermentation. Efforts directed at understanding *T. thermophilus* metabolism revealed that this bacterium uses glycolysis and the TCA cycle to drive carbon flux and bioenergetics.^{202,203} Molecular genetics tools were developed for *Thermus thermophilus*,²⁰⁴ based on its natural competence, which opened up opportunities for it to be examined as a model thermophile. In fact, the relative stability of thermophilic enzymes and early interest in the genus sparked the undertaking of crystallization projects aimed at characterizing recombinant versions of all the identified coding ORFs from *Thermus thermophilus*.^{205–209} The overall goal of such projects was to provide a comprehensive database of structural characteristics that aid in the determination of protein function and domain architecture representing all of the major classes of proteins identified to date.

Enzyme and Metabolic Engineering Efforts with *Thermus*

Although more thermophilic microorganisms have become available, *Thermus* species can be sources of thermostable enzymes for biotechnological applications. For example, enzymes from *Thermus* were included in an *in vitro* pathway that converted glucose into *n*-butanol²¹⁰ and a xylose isomerase from this species was used to enable a recombinant *Saccharomyces cerevisiae* strain to grow on xylose.²¹¹ Although genetics are relatively facile for these organisms, metabolic engineering pursuits have been limited. One of the earliest examples of metabolic engineering of the organism involved the transfer of nitrification genes among two members of the genus,

allowing an aerobic *Thermus* species to grow anaerobically.²¹² Additionally, a few attempts have been made at overexpression of native genes for the purpose of biotechnological applications involving specific enzymes such as DNA polymerase and Mn-dependent catalases.^{213,214} More recently, a strain of *T. thermophilus* HB8 was generated that could co-utilize xylose and glucose at temperatures up to 81°C, with a view towards processing lignocellulose, although this strain could not deconstruct biomass nor ferment the C5/C6 sugars.²¹⁵

CONCLUSION

Although the study of extreme thermophiles has only gained traction in the past few decades, there are numerous metabolic and physiological features that distinguish these organisms from the other major groups of life and justify continued research endeavors. Much of this information has been ascertained via the use of genomics, transcriptomics, and proteomics in conjunction with traditional microbiological/biochemical techniques. Furthermore, this synthesis has led to the development of metabolic and physiological models in extreme thermophiles that are beginning to rival better characterized mesophilic systems. With the advent of next-generation sequencing technologies, it seems likely that previous work will be furthered by large-scale comparative genomics and metagenomics projects; this should further the discovery of novel metabolic features (i.e., enzymes and native biological pathways) with vital importance to our fundamental understanding of biology.

Beyond the scientific merit of studying extreme thermophiles, numerous opportunities exist to utilize these organisms for biotechnological advancement. As previously emphasized, the extreme conditions under which these organisms subsist has led to evolutionarily distinct metabolic and physiological features. In general, thermally stable proteins and heat-tolerant metabolic hosts could provide a major economic benefit to industrial processes. In the case of upstream processes, it may be possible to eliminate or minimize the energy costs associated with cooling or sterilizing bioreactors; while downstream processes may benefit from simple techniques—such as heat pre-treatment—to select for thermophilic enzymes produced recombinantly in mesophilic hosts, eliminating costly purification steps. Additionally, the increase in available genetic systems in these organisms will open many avenues for metabolic engineering. In fact, these organisms could have vital

roles in the future of bioprocessing ranging from sustainable biochemical engineering to specialty chemical production to the deconstruction of inorganic

and organic raw materials and even the recovery of base, precious and strategic metals.

ACKNOWLEDGMENTS

This work was supported in part by the US National Science Foundation (CBET-1264052, CBET-1264053), the BioEnergy Science Center (BESC), a U.S. Department of Energy Bioenergy Research Center supported by the Office of Biological and Environmental Research in the DOE Office of Science, and the US Air Force Office of Sponsored Research (FA9550-13-1-0236). JAC and BMZ acknowledge support from NIH Biotechnology Traineeships (NIH T32GM008776-16). LLL acknowledges support from U.S. NSF Graduate Fellowship. CTS acknowledges support from a U.S. DoEd GAANN Molecular Biotechnology Fellowship.

FURTHER READING

Martin W, Baross J, Kelley D, Russell M. Hydrothermal vents and the origin of life. *Nat Rev Microbiol* 2008, 6:805–814.

REFERENCES

1. Setchell WA. The upper temperature limits of life. *Science* 1903, 17:934–937.
2. Brock TD, Freeze H. *Thermus aquaticus* gen. n. and sp. n., a non-sporulating extreme thermophile. *J Bacteriol* 1969, 98:289–297.
3. Zillig W, Stetter KO, Wunderl S, Schulz W, Priess H, Scholz I. The *Sulfolobus*-*Caldariella* group: taxonomy on the basis of the structure of DNA-dependent RNA polymerases. *Arch Microbiol* 1980, 125: 259–269.
4. Stetter KO, Lauerer G, Thomm M, Neuner A. Isolation of extremely thermophilic sulfate reducers: evidence for a novel branch of archaeobacteria. *Science* 1987, 236:822–824.
5. Huber G, Spinnler C, Gambacorta A, Stetter KO. *Metallosphaera sedula* gen. and sp. nov. represents a new genus of aerobic, metal-mobilizing, thermoacidophilic archaeobacteria. *Syst Appl Microbiol* 1989, 12:38–47.
6. Jannasch HW, Huber R, Belkin S, Stetter KO. *Thermotoga neapolitana* sp. nov. of the extremely thermophilic, eubacterial genus *Thermotoga*. *Arch Microbiol* 1988, 150:103–104.
7. Brock TD, Brock KM, Belly RT, Weiss RL. *Sulfolobus*: a new genus of sulfur-oxidizing bacteria living at low pH and high temperature. *Arch Mikrobiol* 1972, 84:54–68.
8. Bult CA, White O, Olsen GJ, Zhou L, Fleischmann RD, Sutton GG, Blake JA, FitzGerald LM, Clayton RA, Gocayne JD, et al. Complete Genome Sequence of the Methanogenic Archaeon, *Methanococcus jannaschii*. *Science* 1996, 273:1058–1073.
9. Kelnk H-P, Clayton RA, Tomb J-F, White O, Nelson KE, Ketchum KA, Dodson RJ, Gwinn M, Hickey EK, Peterson JD, et al. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* 1997, 390:364–370.
10. Kawarabayasi Y, Sawada M, Horikawa H, Haikawa Y, Hino Y, Yamamoto S, Sekine M, Baba S-I, Kosugi H, Hosoyama A, et al. Complete sequence and gene organization of the genome of a hyperthermophilic archaeobacterium, *Pyrococcus horikoshii* OT3. *DNA Res* 1998, 5:55–76.
11. Zeldes BM, Keller Mw, Loder AJ, Straub CT, Adams MWW, Kelly RM. Extremely thermophilic microorganisms as metabolic engineering platforms for production of fuels and industrial chemicals. *Front Microbiol* 2015, 6:1–17.
12. Albers S-V, Siebers B. The prokaryotes: other major lineages of bacteria and the archaea. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F, eds. *The Family Sulfolobaceae*. Berlin, Heidelberg, Germany. Springer-Verlag; 2006, 323–346. doi:10.1007/0-387-30746-X.
13. Giaveno MA, Urbietta MS, Ulloa JR, González Toril E, Donati ER. Physiologic versatility and growth flexibility as the main characteristics of a novel thermoacidophilic *Acidianus* strain isolated from Copahue Geothermal Area in Argentina. *Microb Ecol* 2013, 65:336–346.
14. Grogan D, Palm P, Zillig W. Isolate B12, which harbours a virus-like element, represents a new species of the archaeobacterial genus *Sulfolobus*, *Sulfolobus shibatae*, sp. nov. *Arch Microbiol* 1990, 154: 594–599.

15. Segerer A, Neuner A, Kristjansson JK, Stetter KO. *Acidianus infernus*, gen. nov., sp. nov. and *Acidianus brierleyi* comb. nov. : facultatively aerobic, extremely acidophilic thermophilic sulfur-metabolizing archaeobacteria. *Int. J. Syst. Bacteriol.* 1986, 36:559–564.
16. Takayanagi S, Kawasaki H, Sugimori K, Yamada T, Sugai A, Ito T, Yamasato K, Shioda M. *Sulfolobus hakenensis* sp. nov., a novel species of acidothermophilic archaeon. *Int J Syst Bacteriol* 1996, 46:377–382.
17. Brierley CL, Brierley JA. A chemoautotrophic and thermophilic microorganism isolated from an acid hot spring. *Can J Microbiol* 1973, 19:183–188.
18. Ding J, Zhang R, Yu Y, Jin D, Liang C, Yi Y, Zhu W, Xia J. A novel acidophilic, thermophilic iron and sulfur-oxidizing archaeon isolated from a hot spring of Tengchong, Yunnan, China. *Braz J Microbiol* 2011, 42:514–525.
19. He ZG, Zhong H, Li Y. *Acidianus tengchongensis* sp. nov., a new species of acidothermophilic archaeon isolated from an acidothermal spring. *Curr Microbiol* 2004, 48:159–163.
20. Fuchs T, Huber H, Teiner K, Burggraf S, Stetter KO. *Metallosphaera prunae*, sp. nov., a novel metal-mobilizing, thermoacidophilic archaeum, isolated from a uranium mine in Germany. *Syst Appl Microbiol* 1995, 18:560–566.
21. Plumb JJ, Haddad CM, Gibson JAE, Franzmann PD. *Acidianus sulfidivorans* sp. nov., an extremely acidophilic, thermophilic archaeon isolated from a solfatar on Lihir Island, Papua New Guinea, and emendation of the genus description. *Int J Syst Evol Microbiol* 2007, 57:1418–1423.
22. Urbietta MS, Rascovan N, Castro C, Revale S, Giaveno MA, Vasquez M, Donati ER. Draft genome sequence of the novel thermoacidophilic archaeon *Acidianus copahuensis* strain ALE1, isolated from the Copahue Volcanic Area in Neuquén, Argentina. *Genome Announc* 2014, 2:e00259–14.
23. Segerer AH, Trincone A, Gahrtz M, Stetter KO. *Stygiolobus azoricus* gen. nov., sp. nov. represents a novel genus of anaerobic, extremely thermoacidophilic archaeobacteria of the order Sulfolobales. *Int J Syst Bacteriol* 1991, 41:495–501.
24. Kurosawa N, Itoh YH, Iwai T, Sugai A, Uda I, Kimura N, Horiuchi T, Itoh T. *Sulfurisphaera ohwakuensis* gen. nov., sp. nov., a novel extremely thermophilic acidophile of the order Sulfolobales. *Int J Syst Bacteriol* 1998, 48:451–456.
25. Grogan DW. Phenotypic characterization of the archaeobacterial genus *Sulfolobus*: Comparison of five wild-type strains. *J Bacteriol* 1989, 171:6710–6719.
26. Zillig W, Kletzin A, Schleper C, Holz I, Janekovic D, Hain J, Lanzendorf M, Kristjansson JK. Screening for Sulfolobales, their plasmids and their viruses in Icelandic solfataras. *Syst Appl Microbiol* 1994, 16:609–628.
27. Dew DW, Buuren CV, McEwan K, Bowker C. Bioleaching of base metal sulphide concentrates: a comparison of high and low temperature bioleaching. *J South Afr Inst Min Metall* 2000, 100:409–414.
28. Norris PR, Burton NP, Clark DA. Mineral sulfide concentrate leaching in high temperature bioreactors. *Miner Eng* 2013, 48:10–19.
29. Wheaton G, Counts J, Mukherjee A, Kruh J, Kelly R. The confluence of heavy metal biooxidation and heavy metal resistance: implications for bioleaching by extreme thermoacidophiles. *Minerals* 2015, 5:397–451.
30. Brierley CL, Brierley JA. Progress in bioleaching: part B: applications of microbial processes by the minerals industries. *Appl Microbiol Biotechnol* 2013, 97:7543–7552.
31. Yoshida N, Nakasato M, Ohmura N, Ando A, Saiki H, Ishii M, Igarashi Y. *Acidianus manzaensis* sp. nov., a novel thermoacidophilic archaeon growing autotrophically by the oxidation of H₂ with the reduction of Fe³⁺. *Curr Microbiol* 2006, 53:406–411.
32. Wächtershäuser G. On the chemistry and evolution of the pioneer organism. *Chem Biodivers* 2007, 4:584–602.
33. Fuchs G. Alternative pathways of carbon dioxide fixation: insights into the early evolution of life? *Annu Rev Microbiol* 2011, 65:631–658.
34. Berg IA. Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl Environ Microbiol* 2011, 77:1925–1936.
35. Bar-Even A, Flamholz A, Noor E, Milo R. Thermodynamic constraints shape the structure of carbon fixation pathways. *Biochim Biophys Acta Bioenergy* 2012, 1817:1646–1659.
36. Berg IA, Kockelkorn D, Buckel W, Fuchs G. A 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway in archaea. *Science* 2007, 318:1782–1786.
37. Hügler M, Krieger RS, Jahn M, Fuchs G. Characterization of acetyl-CoA/propionyl-CoA carboxylase in *Metallosphaera sedula*: carboxylating enzyme in the 3-hydroxypropionate cycle for autotrophic carbon fixation. *Eur J Biochem* 2003, 270:736–744.
38. Estelmann S, Hügler M, Eisenreich W, Werner K, Berg IA, Ramos-Vera WH, Say RF, Kockelkorn D, Gad'on N, Fuchs G. Labeling and enzyme studies of the central carbon metabolism in *Metallosphaera sedula*. *J Bacteriol* 2011, 193:1191–1200.
39. Menendez C, Bauer Z, Huber H, Gad'on N, Stetter KO, Fuchs G. Presence of acetyl coenzyme A (CoA) carboxylase and propionyl-CoA carboxylase in autotrophic crenarchaeota and indication for operation of a 3-hydroxypropionate cycle in autotrophic carbon fixation. *J Bacteriol* 1999, 181:1088–1098.

40. Teufel R, Kung JW, Kockelkorn D, Alber BE, Fuchs G. 3-Hydroxypropionyl-coenzyme A dehydratase and acryloyl-coenzyme A reductase, enzymes of the autotrophic 3-hydroxypropionate/4-hydroxybutyrate cycle in the Sulfolobales. *J Bacteriol* 2009, 191:4572–4581.
41. Ramos-Vera WH, Weiss M, Strittmatter E, Kockelkorn D, Fuchs G. Identification of missing genes and enzymes for autotrophic carbon fixation in *Crenarchaeota*. *J Bacteriol* 2011, 193:1201–1211.
42. Hawkins AS, Han Y, Bennett RK, Adams MWW, Kelly RM. Role of 4-hydroxybutyrate-co a synthetase in the CO₂ fixation cycle in thermoacidophilic archaea. *J Biol Chem* 2013, 288:4012–4022.
43. Alber BE, Kung JW, Fuchs G. 3-Hydroxypropionyl-coenzyme A synthetase from *Metallosphaera sedula*, an enzyme involved in autotrophic CO₂ fixation. *J Bacteriol* 2008, 190:1383–1389.
44. Han Y, Hawkins AS, Adams MWW, Kelly RM. Epimerase (Msed_0639) and mutase (Msed_0638 and Msed_2055) convert (S)-methylmalonyl-coenzyme a (CoA) to succinyl-coA in the *Metallosphaera sedula* 3-hydroxypropionate/4-hydroxybutyrate cycle. *Appl Environ Microbiol* 2012, 78:6194–6202.
45. Kockelkorn D, Fuchs G. Malonic semialdehyde reductase, succinic semialdehyde reductase, and succinyl-coenzyme A reductase from *Metallosphaera sedula*: enzymes of the autotrophic 3-hydroxypropionate/4-hydroxybutyrate cycle in Sulfolobales. *J Bacteriol* 2009, 191:6352–6362.
46. Wery T, Petersen G. Top value added chemicals from biomass volume I — results of screening for potential candidates from sugars and synthesis gas. *U.S. Department of Energy* 2004. doi:10.2172/15008859.
47. Keller MW, Schut GJ, Lipscomb GL, Menon AL, Iwuchukwu IJ, Leuko TT, Thorgersen MP, Nixon WJ, Hawkins AS, Kelly RM, et al. Exploiting microbial hyperthermophilicity to produce an industrial chemical, using hydrogen and carbon dioxide. *Proc Natl Acad Sci USA* 2013, 110:5840–5845.
48. Lian H, Zeldes BM, Lipscomb GL, Hawkins AB, Han Y, Loder AJ, Nishiyama D, Adams MWW, Kelly RM. Ancillary contributions of heterologous biotin protein ligase and carbonic anhydrase for CO₂ incorporation into 3-hydroxypropionate by metabolically engineered *Pyrococcus furiosus*. *Biotechnol Bioeng* 2016, 113:2652–2660. doi:10.1002/bit.26033.
49. Suzuki I. Oxidation of inorganic sulfur compounds: chemical and enzymatic reactions. *Can J Microbiol* 1999, 45:97–105.
50. Kletzin A. Metabolism of inorganic sulfur compounds in Archaea. In: Garrett RA, Klenk H-P, eds. *Archaea*. Malden, MA: Blackwell Publishing Ltd; 2006, 261–274.
51. Blumentals II, Itoh M, Olson GJ, Kelly RM. Role of polysulfides in reduction of elemental sulfur by the hyperthermophilic archaeobacterium *Pyrococcus furiosus*. *Appl Environ Microbiol* 1990, 56:1255–1262.
52. Rabus R, Hansen TA, Widdel F. Dissimilatory sulfate- and sulfur-reducing Prokaryotes. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E, eds. *The Prokaryotes*. New York: Springer; 2006, 659–768.
53. Schauder R, Müller E. Polysulfide as a possible substrate for sulfur-reducing bacteria. *Arch Microbiol* 1993, 160:377–382.
54. Kletzin A. Coupled enzymatic production of sulfite, thiosulfate, and hydrogen sulfide from sulfur: purification and properties of a sulfur oxygenase reductase from the facultatively anaerobic archaeobacterium *Desulfurolobus ambivalens*. *J Bacteriol* 1989, 171:1638–1643.
55. Veith A, Urich T, Seyfarth K, Protze J, Frazão C, Kletzin A. Substrate pathways and mechanisms of inhibition in the sulfur oxygenase reductase of *Acidianus ambivalens*. *Front Microbiol* 2011, 2:1–12.
56. Kletzin A. Oxidation of sulfur and inorganic sulfur compounds in *Acidianus ambivalens*. In: Dahl C, Friedrich CG, eds. *Microbial Sulfur Metabolism*. Berlin/Heidelberg: Springer; 2008, 184–201.
57. Zimmermann P, Laska S, Kletzin A. Two modes of sulfite oxidation in the extremely thermophilic and acidophilic archaeon *Acidianus ambivalens*. *Arch Microbiol* 1999, 172:76–82.
58. Kletzin A. Molecular characterization of the *sor* gene, which encodes the sulfur oxygenase/reductase of the thermoacidophilic archaeum *Desulfurolobus ambivalens*. *J Bacteriol* 1992, 174:5854–5859.
59. Müller FH, Bandejas TM, Urich T, Teixeira M, Gomes CM, Kletzin A. Coupling of the pathway of sulphur oxidation to dioxygen reduction: characterization of a novel membrane-bound thiosulphate:quinone oxidoreductase. *Mol Microbiol* 2004, 53:1147–1160.
60. Purschke WG, Schmidt CL, Petersen A, Schäfer G. The terminal quinol oxidase of the hyperthermophilic archaeon *Acidianus ambivalens* exhibits a novel subunit structure and gene organization. *J Bacteriol* 1997, 179:1344–1353.
61. Anemüller S, Schmidt CL, Pacheco I, Schäfer G, Teixeira M. A cytochrome aa₃-type quinol oxidase from *Desulfurolobus ambivalens*, the most acidophilic archaeon. *FEMS Microbiol Lett* 1994, 117:275–280.
62. Morales P, Gentina J, Morales M, Aroca G, Silva J. Biofiltration of hydrogen sulfide by *Sulfolobus metallicus* at high temperatures. *Water Sci Technol* 2012, 66:1958–1961.
63. Laska S, Lottspeich F, Kletzin A. Membrane-bound hydrogenase and sulfur reductase of the hyperthermophilic and acidophilic archaeon *Acidianus ambivalens*. *Microbiology* 2003, 149:2357–2371.

64. Bathe S, Norris PR. Ferrous iron- and sulfur-induced genes in *Sulfolobus metallicus*. *Appl Environ Microbiol* 2007, 73:2491–2497.
65. Auernik KS, Kelly RM. Identification of components of electron transport chains in the extremely thermoacidophilic crenarchaeon *Metallosphaera sedula* through iron and sulfur compound oxidation transcriptomes. *Appl Environ Microbiol* 2008, 74:7723–7732.
66. Jormakka M, Byrne B, Iwata S. Protonmotive force generation by a redox loop mechanism. *FEBS Lett* 2003, 545:25–30.
67. Barr DW, Ingledew WJ, Norris PR. Respiratory chain components of iron-oxidizing acidophilic bacteria. *FEMS Microbiol Lett* 1990, 70:85–89.
68. Quatrini R, Appia-Ayme C, Denis Y, Jadlicki E, Holmes DS, Bonnefoy V. Extending the models for iron and sulfur oxidation in the extreme acidophile *Acidithiobacillus ferrooxidans*. *BMC Genomics* 2009, 10:394–413.
69. Blake RC, Shute EA, Greenwood MM, Spencer GH, Ingledew WJ. Enzymes of aerobic respiration on iron. *FEMS Microbiol Rev* 1993, 11:9–18.
70. Kozubal MA, Dlakic M, Macur RE, Inskeep WP. Terminal oxidase diversity and function in ‘*Metallosphaera yellowstonensis*’: gene expression and protein modeling suggest mechanisms of Fe(II) oxidation in the Sulfolobales. *Appl Environ Microbiol* 2011, 77:1844–1853.
71. Ilbert M, Bonnefoy V. Insight into the evolution of the iron oxidation pathways. *Biochim Biophys Acta* 2013, 1827:161–175.
72. Clark ME, Batty JD, van Buuren CB, Dew DW, Eamon MA. Biotechnology in minerals processing: Technological breakthroughs creating value. *Hydrometallurgy* 2006, 83:3–9.
73. van Staden PJ, Gericke M, Craven PM. Minerals biotechnology: trends, opportunities and challenges. In: *Hydrometallurgy 2008 : Proceedings of the Sixth International Symposium* 6–13 Society for Mining Metallurgy and Exploration, 2008.
74. Gericke M, Pinches A, van Rooyen J. Bioleaching of a chalcopyrite concentrate using an extremely thermophilic culture. *Int J Miner Process* 2001, 62:243–255.
75. Gericke M, Govender Y, Pinches A. Tank bioleaching of low-grade chalcopyrite concentrates using redox control. *Hydrometallurgy* 2010, 104:414–419.
76. Norris PR, Calvo-Bado L a, Brown CF, Davis-Belmar CS. Ore column leaching with thermophiles: I, copper sulfide ore. *Hydrometallurgy* 2012, 127–128: 62–69.
77. Romano P, Blázquez ML, Alguacil FJ, Muñoz JA, Ballster A, González. Comparative study on the selective chalcopyrite bioleaching of a molybdenite concentrate with mesophilic and thermophilic bacteria. *FEMS Microbiol Lett* 2001, 196:71–75.
78. Abdollahi H, Shafaei SZ, Naoparast M, Manafi Z, Niemelä SI, Tuovinen OH. Mesophilic and thermophilic bioleaching of copper from a chalcopyrite-containing molybdenite concentrate. *Int J Miner Process* 2014, 128:25–32.
79. Takatsugi K, Sasaki K, Hirajima T. Mechanism of the enhancement of bioleaching of copper from enargite by thermophilic iron-oxidizing archaea with the concomitant precipitation of arsenic. *Hydrometallurgy* 2011, 109:90–96.
80. Johnson DB. Biomining-biotechnologies for extracting and recovering metals from ores and waste materials. *Curr Opin Biotechnol* 2014, 30:24–31.
81. Fiala G, Stetter KO. *Pyrococcus furiosus* sp. nov. represents novel genus of marine heterotrophic archaeobacteria growing optimally at 100°C. *Arch Microbiol* 1986, 145:56–61.
82. Kengen SWM, Luesink EJ, Stams AJM, Zehnder AJB. Purification and characterization of an extremely thermostable beta-glucosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Eur J Biochem* 1993, 213:305–312.
83. Gao J, Bauer MW, Shockley KR, Pysz MA, Kelly RM. Growth of hyperthermophilic archaeon *Pyrococcus furiosus* on chitin involves two family 18 chitinases. *Appl Environ Microbiol* 2003, 69:3119.
84. Kreuzer M, Schmutzler K, Waage I, Thomm M, Hausner W. Genetic engineering of *Pyrococcus furiosus* to use chitin as a carbon source. *BMC Biotechnol* 2013, 13:1–10.
85. Snowden LJ, Blumentals II, Kelly RM. Regulation of proteolytic activity in the hyperthermophile *Pyrococcus furiosus*. *Appl Environ Microbiol* 1992, 58:1134–1141.
86. Kengen SWM, de Bok FAM, van Loo N-D, Dijkema C, Stams AJM, de Vos WM. Evidence for the operation of a novel Embden-Meyerhof pathway that involves ADP-dependent kinases during sugar fermentation by *Pyrococcus furiosus*. *J Biol Chem* 1994, 269:17537–17541.
87. Sapra R, Bagramyan K, Adams MWW. A simple energy-conserving system: proton reduction coupled to proton translocation. *Proc Natl Acad Sci USA* 2003, 100:7545–7550.
88. Mai X, Adams MW. Purification and characterization of two reversible and ADP-dependent acetyl coenzyme A synthetases from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Bacteriol* 1996, 178: 5897–5903.
89. Schicho RN, Snowden LJ, Makund S, Park J-B, Adams MWW, Kelly RM. Influence of tungsten on metabolic patterns in *Pyrococcus furiosus*, a hyperthermophilic archaeon. *Arch Microbiol* 1993, 159:380–385.

90. Bryant FO, Adams MW. Characterization of hydrogenase from the hyperthermophilic archaeobacterium, *Pyrococcus furiosus*. *J Biol Chem* 1989, 264: 5070–5079.
91. Schut GJ, Boyd ES, Peters JW, Adams MWW. The modular respiratory complexes involved in hydrogen and sulfur metabolism by heterotrophic hyperthermophilic archaea and their evolutionary implications. *FEMS Microbiol Rev* 2013, 37:182–203.
92. Adams MWW, Holden JF, Menon AL, Schut GJ, Grunden AM, Hou C, Hutchins AM, Jenney Jr. FE, Kim C, Ma K et al. Key role for sulfur in peptide metabolism and in regulation of three hydrogenases in the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Bacteriol* 2001, 183:716–724.
93. Bridger SL, Clarkson Sm, Stirrett K, DeBarry MB, Lipscomb GL, Schut GJ, Westpheling J, Scott RA, Adams MWW. Deletion strains reveal metabolic roles for key elemental sulfur-responsive proteins in *Pyrococcus furiosus*. *J Bacteriol* 2011, 193:6498–6504.
94. Schut GJ, Bridger SL, Adams MWW. Insights into the metabolism of elemental sulfur by the hyperthermophilic archaeon *Pyrococcus furiosus*: characterization of a coenzyme A-dependent NAD(P)H sulfur oxidoreductase. *J Bacteriol* 2007, 189:4431–4441.
95. Lipscomb GL, Keese AM, Cowart DM, Schut GJ, Thomm M, Adams MWW, Scott RA. SurR: A transcriptional activator and repressor controlling hydrogen and elemental sulphur metabolism in *Pyrococcus furiosus*. *Mol Microbiol* 2009, 71:332–349.
96. Yang H, Lipscomb GL, Keese AM, Schut GJ, Thomm M, Adams MWW, Wang BC, Scott RA. SurR regulates hydrogen production in *Pyrococcus furiosus* by a sulfur-dependent redox switch. *Mol Microbiol* 2010, 77:1111–1122.
97. Kengen SWM, Stams AJM. Formation of L-alanine as a reduced end product in carbohydrate fermentation by the hyperthermophilic archaeon *Pyrococcus furiosus*. *Arch Microbiol* 1994, 161:168–175.
98. Lundberg KS, Shoemaker DD, Adams MWW, Short JM, Sorge JA, Mathur EJ. High-fidelity amplification using a thermostable DNA polymerase isolated from *Pyrococcus furiosus*. *Gene* 1991, 108:1–6.
99. Niehaus F, Bertoldo C, Kähler M, Antrankikian G. Extremophiles as a source of novel enzymes for industrial application. *Appl Microbiol Biotechnol* 1999, 51:711–729.
100. Dong GQ, Vieille C, Zeikus JG, Savchenko A, Zeikus JG. Cloning, sequencing, and expression of the gene encoding extracellular alpha-amylase from *Pyrococcus furiosus* and biochemical characterization of the recombinant enzyme. *Appl Environ Microbiol* 1997, 63:3569–3576.
101. Bauer MW, Driskill LE, Kelly RM. An endoglucanase, EglA, from the hyperthermophilic archaeon *Pyrococcus furiosus* hydrolyzes β -1,4 bonds in mixed-linkage (1 \rightarrow 3),(1 \rightarrow 4)- β -D-glucans and cellulose. *J Bacteriol* 1999, 181:284.
102. Bauer MW, Kelly RM. The family 1 beta-glucosidases from *Pyrococcus furiosus* and *Agrobacterium faecalis* share a common catalytic mechanism. *Biochemistry* 1998, 37:17170–17178.
103. Blumentals II, Robinson AS, Kelly RM. Characterization of sodium dodecyl sulfate-resistant proteolytic activity in the hyperthermophilic archaeobacterium *Pyrococcus furiosus*. *Appl Environ Microbiol* 1990, 56:1992–1998.
104. Bauer M, Bauer S, Kelly R. Purification and characterization of a proteasome from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Appl Environ Microbiol* 1997, 63:1160–1164.
105. Farkas J, Stirrett K, Lipscomb GL, Nixon W, Scott RA, Adams MWW, Westpheling J. Recombinogenic properties of *Pyrococcus furiosus* strain COM1 enable rapid selection of targeted mutants. *Appl Environ Microbiol* 2012, 78:4669–4676.
106. Lipscomb GL, Stirrett K, Schut GJ, Yang F, Jenney Jr. FE, Scott RA, Adams MWW, Westpheling J. Natural competence in the hyperthermophilic archaeon *Pyrococcus furiosus* facilitates genetic manipulation: construction of markerless deletions of genes encoding the two cytoplasmic hydrogenases. *Appl Environ Microbiol* 2011, 77:2232–2238.
107. Schut GJ, Lipscomb GL, Nguyen DMN, Kelly RM, Adams MWW. Heterologous production of an energy-conserving carbon monoxide dehydrogenase complex in the hyperthermophile *Pyrococcus furiosus*. *Front Microbiol* 2016, 7:1–9.
108. Hawkins AB, Lian H, Zeldes BM, Loder AJ, Lipscomb GL, Schut GJ, Keller MW, Adams MWW, Kelly RM. Bioprocessing analysis of *Pyrococcus furiosus* strains engineered for CO₂-based 3-hydroxypropionate production. *Biotechnol Bioeng* 2015, 112:1533–1543.
109. Nguyen DMN, Lipscomb GL, Schut GJ, Vaccaro BJ, Basen M, Kelly RM, Adams MWW. Temperature-dependent acetoin production by *Pyrococcus furiosus* is catalyzed by a biosynthetic acetolactate synthase and its deletion improves ethanol production. *Metab Eng* 2016, 34:71–79.
110. Basen M, Sun J, Adams MWW. Engineering a hyperthermophilic archaeon for temperature-dependent product formation. *MBio* 2012, 3:1–8.
111. Keller MW, Lipscomb GL, Loder AJ, Schut GJ, Kelly RM, Adams MWW. A hybrid synthetic pathway for butanol production by a hyperthermophilic microbe. *Metab Eng* 2015, 27:101–106.
112. Blumer-Schuette SE, Brown SD, Sander KB, Bayer EA, Kataeva I, Zurawski JV, Conway JM, Adams MWW, Kelly RM. Thermophilic lignocellulose deconstruction. *FEMS Microbiol Rev* 2014, 38:393–448.

113. VanFossen AL, Ozdemir I, Zelin SL, Kelly RM. Glycoside hydrolase inventory drives plant polysaccharide deconstruction by the extremely thermophilic bacterium *Caldicellulosiruptor saccharolyticus*. *Biotechnol Bioeng* 2011, 108:1559–1569.
114. Zurawski JV, Conway JM, Lee LL, Simpson HJ, Izquierdo JA, Blumer-Schuetz S, Nookaew I, Adams MWW, Kelly RM. Comparative analysis of extremely thermophilic *Caldicellulosiruptor* species reveals common and unique cellular strategies for plant biomass utilization. *Appl Environ Microbiol* 2015, 81:7159–7170.
115. Blumer-Schuetz SE, Giannone RJ, Zurawski JV, Ozdemir I, Ma Q, Yin Y, Xu Y, Kataeva I, Poole II FL, Adams MWW, et al. *Caldicellulosiruptor* core and pangenomes reveal determinants for noncellulosomal thermophilic deconstruction of plant biomass. *J Bacteriol* 2012, 194:4015–4028.
116. Blumer-Schuetz SE, Lewis DL, Kelly RM. Phylogenetic, microbiological, and glycoside hydrolase diversities within the extremely thermophilic, plant biomass-degrading genus *Caldicellulosiruptor*. *Appl Environ Microbiol* 2010, 76:8084–8092.
117. Vanfossen AL, Verhaart MRA, Kengen SMW, Kelly RM. Carbohydrate utilization patterns for the extremely thermophilic bacterium *Caldicellulosiruptor saccharolyticus* reveal broad growth substrate preferences. *Appl Environ Microbiol* 2009, 75:7718–7724.
118. Vanfossen AL, Lewis DL, Nichols JD, Kelly RM. Polysaccharide degradation and synthesis by extremely thermophilic anaerobes. *Ann N Y Acad Sci* 2008, 1125:322–337.
119. Bredholt S, Sonne-Hansen J, Nielsen P, Mathrani IM, Ahring BK. *Caldicellulosiruptor kristjanssonii* sp. nov., a cellulolytic, extremely thermophilic, anaerobic bacterium. *Int J Syst Bacteriol* 1999, 49:991–996.
120. Bielen AAM, Verhaart MRA, van der Oost J, Kengen SWM. Biohydrogen production by the thermophilic bacterium *Caldicellulosiruptor saccharolyticus*: current status and perspectives. *Life (Basel, Switzerland)* 2013, 3:52–85.
121. van de Werken HJG, Verhaart MRA, VanFossen AL, Willquist K, Lewis DL, Nichols JD, Goorissen HP, Mongodin EF, Nelson KE, van Niel EWJ, et al. Hydrogenomics of the extremely thermophilic bacterium *Caldicellulosiruptor saccharolyticus*. *Appl Environ Microbiol* 2008, 74:6720–6729.
122. Zurawski JV, Blumer-Schuetz SE, Conway JM, Kelly RM. The extremely thermophilic genus. In: Zannoni D, De Philippis R, eds. *Advances in Photosynthesis and Respiration, Volume on Microbial Bioenergy: Hydrogen Production*. Netherlands: Springer Science+Business Media; 2014, 177–195. doi:10.1007/978-94-017-8554-9_8.
123. Mladenovska Z, Mathrani IM, Ahring BK. Isolation and characterization of *Caldicellulosiruptor lactoacetivus* sp. nov., an extremely thermophilic, cellulolytic, anaerobic bacterium. *Arch Microbiol* 1995, 163:223–230.
124. Hamilton-Brehm SD, Mosher JL, Vishnivetskaya T, Podar M, Carroll S, Allman S, Phelps TJ, Keller M, Elkins JG. *Caldicellulosiruptor obsidiansis* sp. nov., an anaerobic, extremely thermophilic, cellulolytic bacterium isolated from Obsidian Pool, Yellowstone National Park. *Appl Environ Microbiol* 2010, 76:1014–1020.
125. de Vrije T, Mars AE, Budde AW, Lai MH, Dijkema C, de Waard P, Claassen PAM. Glycolytic pathway and hydrogen yield studies of the extreme thermophile *Caldicellulosiruptor saccharolyticus*. *Appl Microbiol Biotechnol* 2007, 74:1358–1367.
126. Brunecky R, Alahuhta M, Xu Q, Donohoe BS, Crowley MF, Kataeva IA, Yan S-J, Resch MG, Adams MWW, Lunin VV, et al. Revealing nature's cellulase diversity: the digestion mechanism of *Caldicellulosiruptor bescii* CelA. *Science* 2013, 342:1513–1516.
127. Chung D, Young J, Bomble YJ, Vander Wall TA, Groom J, Himmel ME, Westpheling J. Homologous expression of the *Caldicellulosiruptor bescii* CelA reveals that the extracellular protein is glycosylated. *PLoS One* 2015, 10:e0119508.
128. Young J, Chung D, Bomble YJ, Himmel ME, Westpheling J. Deletion of *Caldicellulosiruptor bescii* CelA reveals its crucial role in the deconstruction of lignocellulosic biomass. *Biotechnol Biofuels* 2014, 7:142–149.
129. Ozdemir I, Blumer-Schuetz SE, Kelly RM. S-layer homology domain proteins Csa_c0678 and Csa_c2722 are implicated in plant polysaccharide deconstruction by the extremely thermophilic bacterium *Caldicellulosiruptor saccharolyticus*. *Appl Environ Microbiol* 2012, 78:768–777.
130. Conway JM, Pierce WS, Le JH, Harper GW, Wright JH, Tucker AL, Zurawski JV, Lee LL, Blumer-Schuetz SE, Kelly RM. Multidomain, surface layer-associated glycoside hydrolases contribute to plant polysaccharide degradation by *Caldicellulosiruptor* species. *J Biol Chem* 2016, 291:6732–6747.
131. Blumer-Schuetz SE, Alahuhta M, Conway JM, Lee LL, Zurawski JV, Giannone RJ, Hettich RL, Lunin VV, Himmel ME, Kelly RM. Discrete and structurally unique proteins (tapirins) mediate attachment of extremely thermophilic *Caldicellulosiruptor* species to cellulose. *J Biol Chem* 2015, 290:10645–10656.
132. Basen M, Rhaesa AM, Kataeva I, Pyrbol CJ, Scott IM, Poole FL, Adams MWW. Degradation of high loads of crystalline cellulose and of unpretreated plant biomass by the thermophilic bacterium

- Caldicellulosiruptor bescii*. *Bioresour Technol* 2014, 152:384–392.
133. Bing W, Wang H, Zheng B, Zhang F, Zhu G, Feng Y, Zhang Z. *Caldicellulosiruptor changbaiensis* sp. nov., a cellulolytic and hydrogen-producing bacterium from a hot spring. *Int J Syst Evol Microbiol* 2015, 65:293–297.
 134. Dam P, Kataeva I, Yang S-J, Zhou F, Yin Y, Chou W, Poole II FL, Westpheling J, Hettich R, Giannone R, et al. Insights into plant biomass conversion from the genome of the anaerobic thermophilic bacterium *Caldicellulosiruptor bescii* DSM 6725. *Nucleic Acids Res* 2011, 39:3240–3254.
 135. Isern NG, Xue J, Rao JV, Cort JR, Ahring BK. Novel monosaccharide fermentation products in *Caldicellulosiruptor saccharolyticus* identified using NMR spectroscopy. *Biotechnol Biofuels* 2013, 6:47.
 136. Miroshnichenko ML, Kublanov IV, Kastrikin NA, Tourova TP, Kolganova TV, Birkeland N-K, Bonch-Osmolovskaya EA. *Caldicellulosiruptor kronotskyensis* sp. nov. and *Caldicellulosiruptor hydrothermalis* sp. nov., two extremely thermophilic, cellulolytic, anaerobic bacteria from Kamchatka thermal springs. *Int J Syst Evol Microbiol* 2008, 58:1492–1496.
 137. Bielen AAM, Verhaart MRA, VanFossen AL, Blumer-Schuette SE, Stams AJM, van der Oost J, Kelly RM, Kengen SWM. A thermophile under pressure: transcriptional analysis of the response of *Caldicellulosiruptor saccharolyticus* to different H₂ partial pressures. *Int J Hydrogen Energy* 2013, 38:1837–1849.
 138. Zeidan AA, Van Niel EWJ. Developing a thermophilic hydrogen-producing co-culture for efficient utilization of mixed sugars. *Int J Hydrogen Energy* 2009, 34:4524–4528.
 139. van Niel EWJ, Budde MAW, de Haas GG, van der Wal FJ, Claassen PAM, Stam AJM. Distinctive properties of high hydrogen producing extreme thermophiles, *Caldicellulosiruptor saccharolyticus* and *Thermotoga elfii*. *Int J Hydrogen Energy* 2002, 27:1391–1398.
 140. Willquist K, van Niel EWJ. Lactate formation in *Caldicellulosiruptor saccharolyticus* is regulated by the energy carriers pyrophosphate and ATP. *Metab Eng* 2010, 12:282–290.
 141. de Vrije T, Budde MAW, Lips SJ, Bakker RR, Mars AE, Claassen PAM. Hydrogen production from carrot pulp by the extreme thermophiles *Caldicellulosiruptor saccharolyticus* and *Thermotoga neapolitana*. *Int J Hydrogen Energy* 2010, 35:13206–13213.
 142. Herbel Z, Rákhely G, Bagi Z, Ivanova G, Acs N, Kovács KL. Exploitation of the extremely thermophilic *Caldicellulosiruptor saccharolyticus* in hydrogen and biogas production from biomasses. *Environ Technol* 2010, 31:1017–1024.
 143. Ivanova G, Rákhely G, KOVACS K. Hydrogen production from biopolymers by *Caldicellulosiruptor saccharolyticus* and stabilization of the system by immobilization. *Int J Hydrogen Energy* 2008, 33:6953–6961.
 144. Ivanova G, Rákhely G, Kovács KL. Thermophilic biohydrogen production from energy plants by *Caldicellulosiruptor saccharolyticus* and comparison with related studies. *Int J Hydrogen Energy* 2009, 34:3659–3670.
 145. Kádár Z, de Vrije T, van Noorden GE, Budde MAW, Szengyel Z, Réczey K, Claassen AM. Yields from glucose, xylose, and paper sludge hydrolysate during hydrogen production by the extreme thermophile *Caldicellulosiruptor saccharolyticus*. *Appl Biochem Biotechnol* 2004, 113–116:497–508.
 146. Ljunggren M, Willquist K, Zacchi G, van Niel EW. A kinetic model for quantitative evaluation of the effect of hydrogen and osmolarity on hydrogen production by *Caldicellulosiruptor saccharolyticus*. *Biotechnol Biofuels* 2011, 4:31.
 147. Martinez-Porqueras E, Wechselberger P, Herwig C. Effect of medium composition on biohydrogen production by the extreme thermophilic bacterium *Caldicellulosiruptor saccharolyticus*. *Int J Hydrogen Energy* 2013, 38:11756–11764.
 148. Panagiotopoulos IA, Bakker RR, de Vrije T, Koukios EG, Claassen PAM. Pretreatment of sweet sorghum bagasse for hydrogen production by *Caldicellulosiruptor saccharolyticus*. *Int J Hydrogen Energy* 2010, 35:7738–7747.
 149. Pawar SS, Vongkumpeang T, Grey C, van Niel EW. Biofilm formation by designed co-cultures of *Caldicellulosiruptor* species as a means to improve hydrogen productivity. *Biotechnol Biofuels* 2015, 8:19.
 150. Pawar SS, Nkemka VN, Zeidan AA, Murto M, van Niel EWJ. Biohydrogen production from wheat straw hydrolysate using *Caldicellulosiruptor saccharolyticus* followed by biogas production in a two-step uncoupled process. *Int J Hydrogen Energy* 2013, 38:9121–9130.
 151. Talluri S, Raj SM, Christopher LP. Consolidated bio-processing of untreated switchgrass to hydrogen by the extreme thermophile *Caldicellulosiruptor saccharolyticus* DSM 8903. *Bioresour Technol* 2013, 139:272–279.
 152. Van Niel EWJ, Claassen PAM, Stams AJM. Substrate and product inhibition of hydrogen production by the extreme thermophile, *Caldicellulosiruptor saccharolyticus*. *Biotechnol Bioeng* 2003, 81:255–262.
 153. Van Groenestijn JW, Geelhoed JS, Goorissen HP, Meesters KPM, Stams AJM, Claassen PAM. Performance and population analysis of a non-sterile trickle bed reactor inoculated with *Caldicellulosiruptor saccharolyticus*, a thermophilic hydrogen producer. *Biotechnol Bioeng* 2009, 102:1361–1367.

154. Zeidan AA, van Niel EWJ. A quantitative analysis of hydrogen production efficiency of the extreme thermophile *Caldicellulosiruptor owensensis* OLT. *Int J Hydrogen Energy* 2010, 35:1128–1137.
155. Thauer RK, Jungermann K, Decker K. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* 1977, 41:100–180.
156. Willquist K, Pawar SS, Van Niel EWJ. Reassessment of hydrogen tolerance in *Caldicellulosiruptor saccharolyticus*. *Microb Cell Fact* 2011, 10:111.
157. Peters JW, Miller AF, Jones AK, King PW, Adams MWW. Electron bifurcation. *Curr Opin Chem Biol* 2016, 31:146–152.
158. Farkas J, Chung D, Cha M, Copeland J, Grayeski P, Westpheling J. Improved growth media and culture techniques for genetic analysis and assessment of biomass utilization by *Caldicellulosiruptor bescii*. *J Ind Microbiol Biotechnol* 2013, 40:41–49.
159. Chung DD, Cha M, Farkas J, Westpheling J. Construction of a stable replicating shuttle vector for *Caldicellulosiruptor* species: use for extending genetic methodologies to other members of this genus. *PLoS One* 2013, 8:e62881.
160. Chung D, Farkas J, Huddleston JR, Olivar E, Westpheling J. Methylation by a unique α -class N4-cytosine methyltransferase is required for DNA transformation of *Caldicellulosiruptor bescii* DSM6725. *PLoS One* 2012, 7:e43844.
161. Chung D, Farkas J, Westpheling J. Overcoming restriction as a barrier to DNA transformation in *Caldicellulosiruptor* species results in efficient marker replacement. *Biotechnol Biofuels* 2013, 6:82.
162. Cha M, Chung D, Elkins JG, Guss AM, Westpheling J. Metabolic engineering of *Caldicellulosiruptor bescii* yields increased hydrogen production from lignocellulosic biomass. *Biotechnol Biofuels* 2013, 6:85.
163. Chung D, Cha M, Guss AM, Westpheling J. Direct conversion of plant biomass to ethanol by engineered *Caldicellulosiruptor bescii*. *Proc Natl Acad Sci USA* 2014, 111:8931–8936.
164. Chung D, Cha M, Snyder EN, Elkins JG, Guss AM, Westpheling J. Cellulosic ethanol production via consolidated bioprocessing at 75°C by engineered *Caldicellulosiruptor bescii*. *Biotechnol Biofuels* 2015, 8:163.
165. Cha M, Chung D, Westpheling J. Deletion of a gene cluster for [Ni-Fe] hydrogenase maturation in the anaerobic hyperthermophilic bacterium *Caldicellulosiruptor bescii* identifies its role in hydrogen metabolism. *Appl Microbiol Biotechnol* 2016, 100:1823–1831.
166. Chung D, Pattathil S, Biswal AK, Hahn MG, Mohnen D, Westpheling J. Deletion of a gene cluster encoding pectin degrading enzymes in *Caldicellulosiruptor bescii* reveals an important role for pectin in plant biomass recalcitrance. *Biotechnol Biofuels* 2014, 7:147.
167. Bhandari V, Gupta RS. The phylum Thermotogae. In: Rosenberg E, ed. *The Prokaryotes—Other Major Lineages of Bacteria and the Archaea*. Berlin, Heidelberg, Germany: Springer-Verlag; 2014, 989–1010. doi:10.1007/0-387-30743-5.
168. Belkin S, Wirsén CO, Jannasch HW. A new sulfur-reducing, extremely thermophilic eubacterium from a submarine thermal vent. *Appl Environ Microbiol* 1986, 51:1180–1185.
169. Huber R, Langworthy TA, König H, Thomm M, Woese CR, Sleytr UB, Stetter KO. *Thermotoga maritima* sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. *Arch Microbiol* 1986, 144:324–333.
170. Nelson KE, Clayton RA, Gill SR, Gwinn ML, Dodson RJ, Haft DH, Hickey EK, Peterson JD, Nelson WC, Ketchum KA, et al. Evidence for lateral gene transfer between archaea and bacteria from genome sequence of *Thermotoga maritima*. *Nature* 1999, 399:323–329.
171. Rodionov DA, Kurnasov OV, Stec B, Wang Y, Roberts MF, Osterman AL. Genomic identification and in vitro reconstitution of a complete biosynthetic pathway for the osmolyte di-*myo*-inositol-phosphate. *Proc Natl Acad Sci USA* 2007, 104:4279–4284.
172. Butzin NC, LApierre P, Green AG, Swither KS, Gogarten JP, Noll KM. Reconstructed ancestral *myo*-inositol-3-phosphate synthases indicate that ancestors of the *Thermococcales* and *Thermotoga* species were more thermophilic than their descendants. *PLoS One* 2013, 8:e84300.
173. Ben Hania W, Ghodbane R, Postec A, Brochier-Armanet C, Hamdi M, Fardeau M-L, Ollivier B. Cultivation of the first mesophilic representative ('mesotoga') within the order *Thermotogales*. *Syst Appl Microbiol* 2011, 34:581–585.
174. Rodionova IA, Leyn AS, Burkart MD, Boucher N, Noll KM, Osterman AL, Rodionov DA. Novel inositol catabolic pathway in *Thermotoga maritima*. *Environ Microbiol* 2013, 15:2254–2266.
175. Conners SB, Mongodin EF, Johnson MR, Montero CI, Nelson KE, Kelly RM. Microbial biochemistry, physiology, and biotechnology of hyperthermophilic *Thermotoga* species. *FEMS Microbiol Rev* 2006, 30:872–905.
176. Takahata Y, Nishijima M, Hoaki T, Maruyama T. *Thermotoga petrophila* sp. nov. and *Thermotoga naphthophila* sp. nov., two hyperthermophilic bacteria from the Kubiki oil reservoir in Niigata, Japan. *Int J Syst Evol Microbiol* 2001, 51:1901–1909.
177. Balk MM, Weijma J, Stams AJM. *Thermotoga lettinae* sp. nov., a novel thermophilic, methanol-degrading bacterium isolated from a thermophilic

- anaerobic reactor. *Int J Syst Evol Microbiol* 2002, 52:1361–1368.
178. Ravot G, Magot M, Fardeau M-L, Patel BKC, Prensier G, Egan A, Garcia J-L, Ollivier B. *Thermotoga elfii* sp. nov., a novel thermophilic bacterium from an African oil-producing well. *Int J Syst Bacteriol* 1995, 45:308–314.
179. Jeanthon C, Reysenbach A-L, L'Haridon S, Gambacorta A, Pace NR, Glénat P, Prieur D. *Thermotoga subterranea* sp. nov., a new thermophilic bacterium isolated from a continental oil reservoir. *Arch Microbiol* 1995, 164:91–97.
180. Windberger E, Huber R, Trincone A, Fricke H, Stetter KO. *Thermotoga thermarum* sp. nov. and *Thermotoga neapolitana* occurring in African continental sulfataric springs. *Arch Microbiol* 1989, 151:506–512.
181. Conners SB, Montero CI, Comfort DA, Shockley KR, Johnson MR, Chhabra SR, Kelly RM. An expression-driven approach to the prediction of carbohydrate transport and utilization regulons in the hyperthermophilic bacterium *Thermotoga maritima*. *J Bacteriol* 2005, 187:7267–7282.
182. Chhabra SR, Shockley KR, Conners SB, Scott KE, Wolfinger RD, Kelly RM. Carbohydrate-induced differential gene expression patterns in the hyperthermophilic bacterium *Thermotoga maritima*. *J Biol Chem* 2003, 278:7540–7552.
183. Frock AD, Notey JS, Kelly RM. The genus *Thermotoga*: recent developments. *Environ Technol* 2010, 31:1169–1181.
184. Schröder C, Selig M, Schönheit P. Glucose fermentation to acetate, CO₂ and H₂ in the anaerobic hyperthermophilic eubacterium *Thermotoga maritima*: involvement of the Embden-Meyerhof pathway. *Arch Microbiol* 1994, 161:460–470.
185. Selig M, Xavier KB, Santos H, Schönheit P. Comparative analysis of Embden-Meyerhof and Entner-Doudoroff glycolytic pathways in hyperthermophilic archaea and the bacterium *Thermotoga*. *Arch Microbiol* 1997, 167:217–232.
186. Frock AD, Gray SR, Kelly RM. Hyperthermophilic *Thermotoga* species differ with respect to specific carbohydrate transporters and glycoside hydrolases. *Appl Environ Microbiol* 2012, 78:1978–1986.
187. McCutchen CM, Duffaud GD, Leduc P, Petersen ARH, Tayal A, Khan SA, Kelly RM. Characterization of extremely thermostable enzymatic breakers (a-1,6-galactosidase and B-1,4-mannanase) from the hyperthermophilic bacterium *Thermotoga neapolitana* 5068 for hydrolysis of guar gum. *Biotechnol Bioeng* 1996, 52:332–339.
188. Blumer-Schuette SE, Kataeva I, Westpheling J, Adams MW, Kelly RM. Extremely thermophilic microorganisms for biomass conversion: status and prospects. *Curr Opin Biotechnol* 2008, 19:210–217.
189. Xu H, Han D, Xu Z. Expression of heterologous cellulases in *Thermotoga* sp. Strain RQ2 *Caldicellulosiruptor saccharolyticus*. *Biomed Res Int* 2015, 2015:304–523.
190. Han D, Norris SM, Xu Z. Construction and transformation of a *Thermotoga*-*E. coli* shuttle vector. *BMC Biotechnol* 2012, 12:2.
191. Van Ooteghem SA, Beer SK, Yue PC. Hydrogen production by the thermophilic bacterium, *Thermotoga neapolitana*. *Appl Biochem Biotechnol*. 2002, 98–100:177–189.
192. Van Ooteghem SA, Jones A, Van Der Lelie D, Dong B, Mahajan D. H₂ production and carbon utilization by *Thermotoga neapolitana* under anaerobic and microaerobic growth conditions. *Biotechnol Lett* 2004, 26:1223–1232.
193. Munro SA, Zinder SH, Walker LP. The fermentation stoichiometry of *Thermotoga neapolitana* and influence of temperature, oxygen, and pH on hydrogen production. *Biotechnol Prog* 2009, 25:1035–1042.
194. Schut GJ, Adams MWW. The iron-hydrogenase of *Thermotoga maritima* utilizes ferredoxin and NADH synergistically: a new perspective on anaerobic hydrogen production. *J Bacteriol* 2009, 191:4451–4457.
195. Muralidharan V, Rinker KD, Hirsh IS, Bouwer EJ, Kelly RM. Hydrogen transfer between methanogens and fermentative heterotrophs in hyperthermophilic cocultures. *Biotechnol Bioeng* 1997, 56:268–278.
196. Johnson MR, Conners SB, Montero CI, Chou CJ, Shockley KR, Kelly RM. The *Thermotoga maritima* phenotype is impacted by syntrophic interaction with *Methanococcus jannaschii* in hyperthermophilic coculture. *Appl Environ Microbiol* 2006, 72:811–818.
197. Fardeau M-L, Ollivier B, Patel BKC, Magot M, Thomas P, Rimbault A, Rocchiccioli R, Garcia J-L. *Thermotoga hypogea* sp. nov., a xylanolytic, thermophilic bacterium from an oil-producing well. *Int J Syst Bacteriol* 1997, 47:1013–1019.
198. Ying X, Wang Y, Badiei HR, Karanassios V, Ma K. Purification and characterization of an iron-containing alcohol dehydrogenase in extremely thermophilic bacterium *Thermotoga hypogea*. *Arch Microbiol* 2007, 187:499–510.
199. Eram MS, Wong A, Oduaran E, Ma K. Molecular and biochemical characterization of bifunctional pyruvate decarboxylases and pyruvate ferredoxin oxidoreductases from *Thermotoga maritima* and *Thermotoga hypogea*. *J Biochem* 2015, 158:459–466.
200. Nguyen TAD, Han SJ, Kim JP, Kim MS, Sim SJ. Hydrogen production of the hyperthermophilic eubacterium, *Thermotoga neapolitana* under N₂ sparging condition. *Bioresour Technol* 2010, 101: S38–S41.
201. Oshima T, Imahori K. Description of *Thermus thermophilus* (Yoshida and Oshima) comb. nov., a

- nonsporulating thermophilic bacterium from a Japanese thermal spa. *Int J Syst Bacteriol* 1974, 24:102–112.
202. Lee N-R, Lakshmanan M, Aggarwai S, Song J-W, Karimi IA, Lee D-Y, Park J-B. Genome-scale metabolic network reconstruction and in silico flux analysis of the thermophilic bacterium *Thermus thermophilus* HB27. *Microb Cell Fact* 2014, 13:61–73.
203. Swarup A, Lu J, DeWoody KC, Antoniewicz MR. Metabolic network reconstruction, growth characterization and ^{13}C -metabolic flux analysis of the extremophile *Thermus thermophilus* HB8. *Metab Eng* 2014, 24:173–180.
204. Koyama Y, Hoshino T, Tomizuka N, Furukawa K. Genetic transformation of the extreme thermophile *Thermus thermophilus* and of other *Thermus* spp. *J Bacteriol* 1986, 166:338–340.
205. Sazanov LA, Hinchliffe P. Structure of the hydrophilic domain of respiratory complex I from *Thermus thermophilus*. *Science* 2006, 311:1430–1436.
206. Yokoyama K, Ohkuma S, Taguchi H, Yasunaga T, Wakabayashi T, Yoshida M. V-Type H^+ -ATPase/synthase from a thermophilic eubacterium, *Thermus thermophilus*. *J Biol Chem* 2000, 275:13955–13961.
207. Yusupov MM, Yusupova GZ, Baucom A, Lieberman K, Earnest TN, Cate JHD, Noller HF. Crystal structure of the ribosome at 5.5 Å resolution. *Science* 2001, 292:883–896.
208. Selmer M, Dunham CM, Murphy IV FV, Weixlbaumer A, Petry S, Kelly AC, Weir JR, Ramakrishnan V. Structure of the 70S ribosome complexed with mRNA and tRNA. *Science* 2006, 313:1935–1942.
209. Severinov K. RNA polymerase structure-function: insights into points of transcriptional regulation. *Curr Opin Microbiol* 2000, 3:118–125.
210. Krutsakorn B, Honda K, Ye X, Imagawa T, Bei X, Okano K, Ohtake H. In vitro production of *n*-butanol from glucose. *Metab Eng* 2013, 20:84–91.
211. Karhumaa K, Hahn-Hägerdal B, Gorwa-Grauslund MF. Investigation of limiting metabolic steps in the utilization of xylose by recombinant *Saccharomyces cerevisiae* using metabolic engineering. *Yeast* 2005, 22:359–368.
212. Ramírez-Arcos S, Fernández-Herrero LA, Marín I, Berenguer J. Anaerobic growth, a property horizontally transferred by an Hfr-like mechanism among extreme thermophiles. *J Bacteriol* 1998, 180:3137–3143.
213. Moreno R, Haro A, Castellanos A, Berenguer J. High-level overproduction of His-tagged Tth DNA polymerase in *Thermus thermophilus*. *Appl Environ Microbiol* 2005, 71:591–593.
214. Hidalgo A, Betancor L, Moreno R, Zafra O, Cava F, Fernández-Laufuente R, Guisán JM, Berenguer J. *Thermus thermophilus* as a cell factory for the production of a thermophilic Mn-dependent catalase which fails to be synthesized in an active form in *Escherichia coli*. *Appl Environ Microbiol* 2004, 70:3839–3844.
215. Cordova LT, Lu J, Cipolla RM, Sandoval NR, Long CP, Antoniewicz MR. Co-utilization of glucose and xylose by evolved *Thermus thermophilus* LC113 strain elucidated by ^{13}C metabolic flux analysis and whole genome sequencing. *Metab Eng* 2016, 37:63–71.